arXiv:1204.5566v1 [math-ph] 25 Apr 2012

Deformation Expression for Elements of Algebras (VI)
–Vacuum representation of Heisenberg algebra–

Hideki Omori[*] Yoshiaki Maeda[†] Naoya Miyazaki[‡]
Tokyo University of Science Keio University Keio University

Akira Yoshioka [§]
Tokyo University of Science

Contents

1 Introduction **2**

2 μ-regulated algebra **4**
2.1 Abstract vacuums 7

3 Extended Weyl algebra **9**
3.1 Extension of products 11
3.1.1 Several remarks on the notation $:A_*:_{_K}$ 11
3.2 Star-exponential functions 12
3.2.1 The case $\rho \in \mathcal{H}_{2m}[\nu]$ 14
3.2.2 Heisenberg algebra embedded in Weyl algebra 15
3.3 $*$-exponentials of quadratic forms 16
3.3.1 Abstract vacuums on several variables 17
3.4 Reductions vs Restriction to the Energy surface 19
3.4.1 Complex vacuum representation 20
3.4.2 Localizations joining divisors 21

4 The case embedded Heisenberg algebra **21**
4.1 The star-exponential functions $e_*^{it\tau}{*}F$, $F{*}e_*^{-it\tau}$ 24
4.1.1 Solutions for the initial value $e^{a\frac{1}{i\hbar}x_0y_0}$ 28

---

[*] Department of Mathematics, Faculty of Sciences and Technology, Tokyo University of Science, 2641, Noda, Chiba, 278-8510, Japan, email: omori@ma.noda.tus.ac.jp

[†] Department of Mathematics, Faculty of Science and Technology, Keio University, 3-14-1, Hiyoshi, Yokohama,223-8522, Japan, email: maeda@math.keio.ac.jp

[‡] Department of Mathematics, Faculty of Economics, Keio University, 4-1-1, Hiyoshi, Yokohama, 223-8521, Japan, email: miyazaki@hc.cc.keio.ac.jp

[§] Department of Mathematics, Faculty of Science, Tokyo University of Science, 1-3, Kagurazaka, Tokyo, 102-8601, Japan, email: yoshioka@rs.kagu.tus.ac.jp

4.1.2 The case that initial date is $\overline{\varpi}_{00}$ 30
4.1.3 Solution for the initial value 1 31

5 Berezin algebra embedded in the extended Weyl algebra 33
5.1 Algebra on the unit disk D. 34
5.2 Embeddings into the extended Weyl algebra 37

Keywords: Weyl algebra, Heisenberg algebra, Contact Weyl algebra, Vacuum representation, Time-energy uncertainty.
Mathematics Subject Classification(2000): Primary 53D55, Secondary 53D17, 53D10

1 Introduction

The **Weyl algebra** $(W_{2m}[\hbar]; *)$ is the algebra generated by $\boldsymbol{u}$=$(u_1, \cdots, u_m, v_1, \cdots, v_m)$ over $\mathbb{C}$ with the fundamental commutation relation $[u_i, v_j] = -i\hbar\delta_{ij}$, where $\hbar$ is a positive constant. The Heisenberg algebra $(\mathcal{H}_{2m}[\nu]; *)$ is the algebra given by regarding the scalar parameter $\hbar$ in the Weyl algebra $W_{2m}[\hbar]$ to be a generator ν which commutes with all others. The difference of those two algebras is seen in the next

Proposition 1.1 *There is no nontrivial two-sided ideal of the Weyl algebra* $(W_{2m}[\hbar]; *)$*. On the other hand, the Heisenberg algebra* $(\mathcal{H}_{2m}[\nu]; *)$ *has two-sided ideals corresponding to points of* $\mathbb{C}^{2m}$.

Proof is easy by observing the following: Suppose ψ is a homomorphism of an algebra into $\mathbb{C}$, and suppose $[x, y]_* = z$, then $\psi(z) = 0$. It follows that there is no nontrivial two-sided ideal of the Weyl algebra $W_{2m}[\hbar]$.

On the other hand $\nu{*}\mathcal{H}_{2m}[\nu]$ is a two-sided ideal of $\mathcal{H}_{2m}[\nu]$ such that quotient algebra is the usual commutative polynomial ring $\mathbb{C}[\boldsymbol{u}]$. It is easy to see that for every $\boldsymbol{a} \in \mathbb{C}^{2m}$, the two-sided ideal of $\mathbb{C}[\boldsymbol{u}]$ generated by $\boldsymbol{u}{-}\boldsymbol{a}$ is pull back to give an nontrivial two-sided ideal of $\mathcal{H}_{2m}[\nu]$. $\square$

Thus, the Heisenberg algebra may be treated in $(\mathit{Hol}(\mathbb{C}^{2m+1}), *)$. These are seen in [12], pp195-200, pp300-305.

However ν in the Heisenberg algebra is often regarded as an invertible element. In precisely, this means one may join the inverse ν^{-1} to $\mathcal{H}_{2m}[\nu]$. In this case, the extended algebra $\mathcal{H}_{2m}[\nu, \nu^{-1}]$ turns out to be isomorphic to $W_{2m}[\hbar]$. In spite of this, there is big difference in the notion of automorphisms.

Consider a one parameter transformation $E_t : \mathcal{H}_{2m}[\nu] \to \mathcal{H}_{2m}[\nu]$ by setting on generators as

$$E_t(\nu) = e^{2t}\nu, \quad E_t(u_i) = e^t u_i, \quad E_t(v_i) = e^t v_i, \quad i = 1, \ldots, m, \quad t \in \mathbb{R}.$$

E_t extends naturally to an automorphism of the Heisenberg algebra

$$E_t : \mathcal{H}_{2m}[\nu] \to \mathcal{H}_{2m}[\nu], \tag{1}$$

but this is not an automorphism of the Weyl algebra $W_{2m}[\hbar]$, as $\hbar$ is not fixed. We refer E_t to an **expansive automorphism**.

Consider the infinitesimal generator $D = \frac{d}{dt}E_t\Big|_{t=0}$ of (1). Obviously,

$$D(\nu) = 2\nu, \quad D(u_i) = u_i, \quad D(v_i) = v_i, \quad i = 1, \dots, m.$$

Suppose now that D is given by $\frac{1}{\nu}\mathrm{ad}(\frac{1}{i}\tau)$ in the ordinary shape of Hamiltonian mechanics using a certain (virtual) Hamiltonian τ.

We join the virtual element τ to the Heisenberg algebra $\mathcal{H}_{2m}[\nu]$ by setting

$$[i\tau, \nu] = -2\nu^2, \quad [i\tau, u_i] = -\nu{*}u_i, \quad i = 1, \dots, 2m. \tag{2}$$

Though ν^{-1} is not an element of $\mathcal{H}_{2m}[\nu]$, we note $[\nu^{-1}, \tau]{=}2i$.

In Heisenberg algebra, ν^{-1} is often viewed as the (independent universal) "energy variable". Then its canonical conjugate τ may be viewed as the "universal time" which might be fixed by the negotiation among the allover universe just like the Greenwich mean time.

However, the existence of such a canonical conjugate element is very controversially relating to the **time-energy uncertainty**, as one may see in [5]. From a view point of differential geometry, the system (2) is obtained as the deformation quantization of a local normal form of a contact structure. (See Proposition 2.1.)

In this note we are interested in the algebra consisting of E_t-invariant elements. Note that in quantum mechanics there is no strict notion of restricting variables, while it is trivial in differential geometry, where everything is considered on a patchwork of local coordinate system. Thus, we regard this as the algebra of (non-commutative) unit sphere $S^{2m-1}{=}\{\sum_{k=1}^{m}(u_k^2{+}v_k^2){=}1\}$. This algebra is generated by

$$\rho^{-2}{*}\nu,\ \rho^{-1}{*}u_1, \cdots, \rho^{-1}{*}u_m,\ \rho^{-1}{*}v_1, \cdots, \rho^{-1}{*}v_m \tag{3}$$

where ρ is an element satisfying $E_t\rho{=}e^t\rho$. Obviously, ρ^{-1} is *not* an element of $\mathcal{H}_{2m}[\nu]$. Hence we have to join such an element to the algebra $\mathcal{H}_{2m}[\nu]$ to regulate the system, but the generated algebra depends how the commutation relations $[\rho^{-2}, \nu]$, $[\rho^{-1}, u_i]$, $[\rho^{-1}, v_j]$ are defined. If these are 0, then the generated algebra is isomorphic to the Heisenberg algebra $\mathcal{H}_{2m}[\rho^{-2}{*}\nu]$, hence this procedure does not make the restriction of variables.

The case that $\rho_*^2{=}\sum_{i=1}^{m}(u_i^2{+}v_i^2)$ may be the most familiar case. In physics, $i\nu$ and the transcendental element $\sqrt[*]{\sum_{i=1}^{m}(u_i^2{+}v_i^2)}$ are often treated as a selfadjoint semi-bounded operators. This implies that τ cannot be a self-ajoint operator. This is because as follows:

If it is possible, there must be a one parameter group of unitary operators $e_*^{it\tau}$, $t \in \mathbb{R}$. Suppose $e_*^{it\tau}$, $t \in \mathbb{R}$, exists by *itself* and $A_t{=}e_*^{it\tau}{*}\nu^{-1}{*}e_*^{-it\tau}$ is a welldefined real analytic group action in t. We see that A_t must satisfy the equation

$$\frac{d}{dt}A_t{=}\mathrm{ad}(i\tau)A_t, \quad A_0{=}\nu^{-1}.$$

Hence the uniqueness of the real analytic solution and the identity $[i\tau, \nu^{-1}]{=}2$ give

$$e_*^{it\tau}{*}\nu^{-1}{*}e_*^{-it\tau}{=}e^{t\mathrm{ad}(i\tau)}\nu^{-1}{=}\nu^{-1}{+}2t{=}\nu^{-1}(1{+}2\nu t).$$

It follows $\nu^{-1}{+}2t$ is invertible for every $t{\in}\mathbb{R}$. This against the assumption ν is self-adjoint.

On the other hand, as $\rho_*^{-2}{*}\nu$ is E_t-invariant, we have the identity $[\tau, \rho_*^{-2}{*}\nu]{=}0$, which easily yields $[\tau, \rho_*^{-2}]{*}\nu{-}\rho_*^{-2}2i{=}0$, hence $[i\tau, \rho_*^{-2}]{=}2\nu{*}\rho_*^{-2}$, $[i\tau, [i\tau, \rho_*^{-2}]]{=}0, \cdots$. Thus, the real analytic solution of the equation

$$\frac{d}{ds}e_*^{is\tau}{*}\rho_*^{-2}{*}e_*^{-is\tau}{=}\mathrm{ad}(i\tau)e_*^{is\tau}{*}\rho_*^{-2}{*}e_*^{-is\tau}$$

is $e_*^{is\tau}{*}\rho_*^{-2}{*}e_*^{-is\tau}{=}(1{+}2\nu s){*}\rho_*^{-2}$. It follows $\rho_*^{-2}{*}e_*^{-is\tau}{*}\rho_*^{2}{=}(1{+}2\nu s)e_*^{-is\tau}$. That is, $e_*^{-is\tau}$ is an eigenvector of $\mathrm{Ad}(\rho^{-2})$. Hence in the situation that $\mathrm{Ad}(\rho_*^{-2})$ has discrete eigenvectors, we have to think $e_*^{-is\tau}$ forms a discrete set. This may not be realistic.

These observation suggest that $e_*^{it\tau}$ cannot exist by *itself*. That is, the equation

$$\frac{d}{dt}F_t{=}i\tau{*}F_t, \quad F_0{=}f$$

may not be solved for the initial condition 1 or even if it exists solutions may not be unique.

On the other hand in the theory of deformation quantization (cf.[3]), operator representations are required only in the final stage. In the beginning, we have only to extend $W_{2m}[\hbar]$ to a topological two-sided $W_{2m}[\hbar]$-module. In an extended system, one may define $\sqrt[*]{\sum_{i=1}^{m}(u_i^2{+}v_i^2)}$ without using operator representations. One has no need to care about positivity or self-adjointness, as there are no such notions. We are now free from the operator theory.

In this note, we propose an approach to the problem of the time- energy uncertainty from a little more differential geometrical view point together with the notion of abstract vacuums in the next section.

2 μ-regulated algebra

We begin with the notion which will be convenient to understand the motivation of the theory of deformation quantization (cf.[3]), and systems appeared by the restriction of the reduction procedures. Although it is not directly relevant to this note, μ-regulated algebras are also very convenient to treat the calculus of pseudo-differential operators on manifolds (cf.[12]). These will be discussed in forthcoming notes.

Let $\mathcal{A}$ be a topological algebra with 1 over $\mathbb{C}$, which may not be a complete topological space. We denote the product in $\mathcal{A}$ by $a*b$. $\mathcal{A}$ is called a *μ-regulated algebra*, if there is an element $\mu\in\mathcal{A}$, called a *regulator*, such that $(\mathcal{A}, \mu)$ satisfies the following:

(A.1) $[\mu, \mathcal{A}]\subset\mu{*}\mathcal{A}{*}\mu$.

(A.2) $[\mathcal{A}, \mathcal{A}]\subset\mu{*}\mathcal{A}$. ($\mathcal{A}$ is abelian modulo μ.)

(A.3) There is a closed subspace B such that $\mathcal{A}=B\oplus\mu{*}\mathcal{A}$ (topological direct sum).

(A.4) Mappings $\mu{*}:\mathcal{A}\to\mu{*}\mathcal{A}$, ${*}\mu:\mathcal{A}\to\mathcal{A}{*}\mu$ defined by $a\to\mu*a$, $a\to a*\mu$ respectively are linear isomorphisms.

(A.1) shows $a{*}\mu{=}\mu{*}a{+}\mu{*}^{\exists}b{*}\mu{=}\mu{*}(a{+}b{*}\mu)$, hence $\mu{*}\mathcal{A}\subset\mathcal{A}{*}\mu\subset\mu{*}\mathcal{A}$. Thus $\mu{*}\mathcal{A}{=}\mathcal{A}{*}\mu$. It is a closed subspace, and hence $\mu{*}\mathcal{A}$ is a closed two-sided ideal of $\mathcal{A}$. (A.2) shows the factor space $B=\mathcal{A}/\mu{*}\mathcal{A}$ is a topological commutative algebra. Note that complementary space B is not unique in general, but the quotient algebras are mutually isomorphic commutative algebras.

By the properties (A.3), (A.4), $\mathcal{A}$ is decomposed for every N into

$$\mathcal{A}=B\oplus\mu{*}B\oplus\cdots\oplus\mu^{N-1}{*}B\oplus\mu^{N}{*}\mathcal{A}. \tag{4}$$

We set $\mathcal{A}^{-\infty}{=}\bigcap_k\mu^k{*}\mathcal{A}$. Then, $\mu\notin\mathcal{A}^{-\infty}$. If $\mathcal{A}^{-\infty}{=}\{0\}$, $\mathcal{A}$ is said to be *analytic* or *formal*.

For every $a,b\in B$, the decomposition (4) gives

$$a*b\sim\sum\nolimits_{k\geq 0}\mu^k*\pi_k(a,b),\quad \pi_k(a,b)\in B. \tag{5}$$

We see $\pi_0(a,b)=a{\cdot}b$ is an associative commutative product, and the skew part π_1^- of π_1 gives a biderivation of $B\times B$ into B. $\pi_1^-(a,b)$ is denoted by $\{a,b\}$ and often called as Poisson bracket product.

The property (A.4) permits us to join the inverse μ^{-1} to the algebra $\mathcal{A}$. By setting $\mu{*}\mu^{-1}{=}\mu^{-1}{*}\mu{=}1$ and $[\mu^{-1},a]{=}-\mu^{-1}{*}[\mu,a]{*}\mu^{-1}$, $\mathrm{ad}(i\mu^{-1})$ gives a derivation of $\mathcal{A}$, and it is decomposed as

$$\mathrm{ad}(i\mu^{-1})(a)=\xi_0(a)+\mu{*}\xi_1(a){+}\cdots{+}\mu^k{*}\xi_k(a){+}\cdots,\quad a\in B \tag{6}$$

where ξ_0 is a derivation of $(B,\cdot)$. Hence ξ_0 is viewed as a vector field, which is called the *characteristic vector field*. The parameter of its integral curves is often viewed as the "time" in differential geometry. Similarly, $(A.2)$ shows that $\mathrm{ad}(a{*}\mu^{-1})$ is an outer derivation of $\mathcal{A}$ for every $a{\in}\mathcal{A}$, and $(\mathcal{A}{*}\mu^{-1},[\ ,\ ])$ forms a Lie algebra over $\mathbb{C}$ containing $(\mathcal{A},[\ ,\ ])$ as a Lie ideal. The quotient Lie algebra $\mathcal{A}{*}\mu^{-1}/\mathcal{A}$ is called the Jacobi algebra whose bracket product is given by

$$\{f,g\}_c{=}f\xi_0(g){-}g\xi_0(f){+}\{f,g\},\quad f,g\in B.$$

The enveloping associative algebra of $(\mathcal{A}{*}\mu^{-1},[\ ,\ ])$ will be denoted by $\mathcal{A}[\mu^{-1}]$. This is often more important than $(\mathcal{A},*)$ itself or the Lie algebra $(\mathcal{A}{*}\mu^{-1},[\ ,\ ])$. As elements of $\mathcal{A}{*}\mu^{-1}$ are represented by unbounded operators in general, it seems that the complex Lie algebra $(\mathcal{A}{*}\mu^{-1},[\ ,\ ])$ cannot be a Lie algebra of a certain "Lie group" (cf.[12] pp.167-169).

Let $B_0{=}\{f{\in}B;\xi_0(f){=}0\}$. B_0 is a closed subalgebra of $(B,\cdot)$. There may be the case $B_0{=}B$, i.e. $\xi_0{=}0$. If this is the case, then $[\mu^{-1},B]\subset\mu{*}\mathcal{A}$ and π_1^- satisfies the Jacobi identity, that is, (B,π_1^-) is a Lie algebra. If $\xi_0{=}0$ and π_1^- is nondegenerate on the space B, then $(B,\cdot,\pi_1^-,\mu)$ is called an **abstract symplectic structure**. If $\xi_0{\neq}0$ and π_1^- is nondegenerate on the space B_0, then $(B,\cdot,\xi_0,\pi_1^-)$ is called **an abstract contact structure**.

In this context, Heisenberg algebra is often regarded as a subalgebra of an abstract contact algebra or a quantized contact algebra. In this case, Heisenberg algebra appears as $\mathcal{A}_0{=}\{a{\in}\mathcal{A};[\mu^{-1},a]=0\}$.

Although the choice of the complementary space B is not unique, but it depends on the expression parameters mentioned below, it is natural to think that all differential geometrical structures defined on $(B,\cdot)$ should be expressed by notions involved in the first few terms of the μ-series. This gives indeed the motivation of "deformation quantization". If it is possible, then such a geometrical structure is called to be *deformation quantizable.*

If this is the sole question, one has only to make a μ-regulated algebra where μ-series are formal power series, and then the quantization problems would have almost been settled. Namely *every Poisson algebra is deformation quantizable* (cf. [11]).

However, it is natural to think that the theory via formal μ is not a true physics but only a probe toward the non-formal theory.

In differential geometry, $(B, \cdot)$ is often assumed to be a commutative function algebra on a *phase* manifold M, and μ^{-1} is a real valued function on M, which may be called *hamiltonian.*

As $\xi_0(\mu) = 0$, the equality $\mu^{-1}{=}s$ may be regarded as the *energy surface* of level s, and μ^{-1} may be regarded as a local coordinate function on a neighborhood of the surface. A suitable local assumption permits us to consider a local coordinate function τ such that $\xi_0 = \partial_\tau$, hence $\partial_\tau B_0{=}0$ and $\xi_0(\tau) = 1$. Hence in differential geometry, a contact structure appears on the energy surface and τ is involved as a coordinate function of the configuration space (space-time).

In functional analysis, the eigenspace decomposition $\sum_{s\in I} \oplus \mathcal{A}_s$ of $\mathrm{ad}(\mu^{-1}) : \mathcal{A} \to \mathcal{A}$ is treated under suitable assumptions such as semi-boundedness or positivity. Let $\{a_s; s{\in}I\}$ be a family of eigenvectors, i.e. $[\mu^{-1}, a_s] = sa_s$. Then, roughly speaking, we have

$$[\mu^{-1}, \log a_s] = sa_s/a_s = s.$$

Hence, if $\tau_s{=}\log a_s$ exists, then the family $\{\tau_s; s{\in}I\}$ may be viewed as the canonical conjugate variable of $\mu^{-1}{=}\{\mu^{-1}; s{\in}I\}$ by regarding every $f{\in}\mathcal{A}$ as a constant family $\{f; s{\in}I\}$.

A μ-regulated algebra with an abstract contact structure is called a **contact Weyl algebra**. Such an algebra often appears in the shape $\mathcal{A}_0[\tau]$. In general, there is a serious unrepairable gap between these two view points. Main reason of this gap is that local structures are leading target in differential geometry, but the global structures are concerned in analysis. One of the reason is that the method of obtaining *normal form* is given only in the formal level, as it will be mentioned below.

Localization theorem. If a μ-regulated algebra $(\mathcal{A}, \mu)$ is defined by setting a $*$-product on a function space of a manifold M, then by choosing a suitable complementary subspace B, all ξ_k and π_k become respectively differential operators and bidifferential operators. This is called the localization theorem proved in [12]. Note that $\mathcal{A}/\mathcal{A}^{-\infty}$ is a (formal) μ-regulated algebra. The localization theorem shows that $\mathcal{A}/\mathcal{A}^{-\infty}$ can be restricted on every open subset U of M by using the restrictions of π_k and ξ_k on U.

Quantum Darboux theorem (cf.[12]). If the abstract symplectic structure is involved further, then the localization theorem and stepwise repairing of coordinate functions show that there is an open neighborhood U_p of $p{\in}M$ on which $\mathcal{A}/\mathcal{A}^{-\infty}$ is expressed as a Weyl algebra. (See the next section for the definition of the Weyl algebra.)

If such a localization is obtained by joining "divisors", then one may not loose the information about $\mathcal{A}^{-\infty}$. Note also that the μ-regulated algebra $\widetilde{\mathcal{A}}$ reconstructed by localized generator system must have $\widetilde{\mathcal{A}}^{-\infty}$ part. That is, localized system might not be formal. Now it arises a big, basic question how $\widetilde{\mathcal{A}}$ relates to the original $\mathcal{A}$. A system in a laboratory must be a certain localization of the system of the whole universe.

The next Proposition shows that (2) is a standard local coordinate system.

Proposition 2.1 *Suppose $(B,\cdot)$ is the space of C^∞ functions on a manifold M and μ is given as a smooth function on M. Then, there is a local coordinate system with the property* (2) *on a tubular neighborhood of a compact fragment of an orbit of ξ_0.*

Proof (Sketch) As $[\mu^{-1},\mu]{=}0$ gives $\xi_0(\mu)=0$, every orbit of ξ_0 is sitting on a subspace $\mu{=}s$(constant). One may assume $\mu{=}s$ defines a manifold M_s of codimension one. Consider a compact fragment $c(\tau)$ of an orbit of ξ_0. We assume $\xi_0(\tau)=1$, and we take a system of local coordinate $s,u_1,\cdots,u_{m-1}$, $v_1,\cdots,v_{m-1}$ on a disk transversal to $c(*)$. It is clear that $\xi_0(s)=0$ and one may assume $\xi_0(u_i)=0$ and $\xi_0(v_i)=0$ for $i=1\sim m-1$. As π_1^- is nondegenerate, classical Darboux theorem shows that one may assume $\pi_1^-(u_i,v_j){=}\delta_{ij}$.

By the localization theorem, the original $*$-product can be restricted to the functions on this open neighborhood. Hence, we have $\mu{=}s$ and

$$[\mu^{-1},u_k]{=}0,\quad mod(\mu),\quad [\mu^{-1},v_k]{=}0,\quad mod(\mu),\quad [u_i,v_j]=\mu i\delta_{ij}\quad mod(\mu^2)$$

On the other hand, we denote

$$[\begin{bmatrix}\boldsymbol{u}\\ \boldsymbol{v}\end{bmatrix},({}^t\!\boldsymbol{u},{}^t\!\boldsymbol{v})]=2i\mu^2\begin{bmatrix}0&-I\\ I&0\end{bmatrix},\quad \mathrm{ad}(\tau)\begin{bmatrix}\boldsymbol{u}\\ \boldsymbol{v}\end{bmatrix}{=}A\begin{bmatrix}\boldsymbol{u}\\ \boldsymbol{v}\end{bmatrix},\quad \mathrm{ad}(\tau)({}^t\!\boldsymbol{u},{}^t\!\boldsymbol{v})=({}^t\!\boldsymbol{u},{}^t\!\boldsymbol{v}){}^tA.$$

The identity

$$2i\mu^2\delta_{ij}{=}[\tau,\mu\delta_{ij}]=[\tau,[u_i,v_j]]=[[\tau,u_i],v_j]{+}[u_i,[\tau,v_j]]$$

together with $[\tau,[u_i,u_j]]=[\tau,[v_i,v_j]]=0$ gives $2i\mu^2J=AJ{+}J^tA$. It follows

$$(A{-}i\mu)J{+}J^t(A{-}i\mu)=0,\quad i.e.\ (A{-}i\mu)J\ \text{ is a symmetric matrix.}$$

By a suitable linear change of coordinate one may set that

$$[\tau,u_k]{=}a_k\mu{*}u_k{+}\cdots,\quad [\tau,v_k]{=}b_k\mu{*}v_k{+}\cdots,$$

and one may assume $a_k=b_k=i$ after a further tuning. Suitable change of coordinate by a standard linearization procedure gives that $[\tau,u_i]{=}i\mu{*}u_i$, $[\tau,v_i]{=}i\mu{*}v_i$. Embed this system into a symplectic manifold, and the quantum Darboux theorem gives the desired result. $\square$

2.1 Abstract vacuums

In physics, the vacuum may be defined as the lowest eigenstate of energy, but in our situation $\mathrm{ad}(\mu^{-1})$ may not be semi-bounded. Here, we propose a little different notion of vacuums. We call an idempotent element $\varpi{\in}\mathcal{A}$ an **abstract vacuum** if

$$\varpi{*}\varpi{=}\varpi,\quad \varpi{*}\mathcal{A}{*}\varpi{=}\mathbb{C}{*}\varpi. \tag{7}$$

Such an element is not contained in the Weyl algebra, but there are a lot of such elements in a transcendentally extended Weyl algebra. If ϖ is an abstract vacuum and $\varpi{*}g{*}\varpi{=}\lambda_g\varpi$, $\lambda_g\in\mathbb{C}\setminus\{0\}$, then $\lambda_g^{-1}g{*}\varpi$ is an abstract vacuum.

It is remarkable that an abstract vacuum exists only in $\mathcal{A}^{-\infty}$, i.e $\mathcal{A}$ non-formal . This is proved as follows: If $\varpi{*}\mu{*}\varpi{=}c\varpi$, $c\neq 0$, then $\frac{1}{c}\mu{*}\varpi{\in}\mu{*}\mathcal{A}$ is an abstract vacuum. As this is idempotent, we see $\frac{1}{c}\mu{*}\varpi{\in}\mathcal{A}^{-\infty}$ and hence $\varpi{\in}\mathcal{A}^{-\infty}$. If $c{=}0$, then $(A.1)$ gives $0{=}\varpi{*}\mu{*}\varpi{=}\mu{*}\varpi{+}\mu{*}a{*}\mu{*}\varpi$. Hence, $\mu{*}\varpi{\in}\mu^2{*}\mathcal{A}$, and then $\varpi{\in}\mu{*}\mathcal{A}$. It follows $\varpi{\in}\mathcal{A}^{-\infty}$, and $\mathcal{A}{*}\varpi{*}\mathcal{A}\subset\mathcal{A}^{-\infty}$.

If $\mathcal{A}$ is generated by $1, x_1, \cdots, x_n$, and $x_i{*}\varpi{=}0$ or $\varpi{*}x_i{=}0$ holds for every i, then ϖ is an abstract vacuum. We call such an abstract vacuum a **standard abstract vacuum**. (Note that there is a non-standard abstract vacuum, called a pseudo-vacuum in [19].)

Sometimes, it is natural to assume

$$\mu{*}\varpi{=}c\varpi, \quad c>0. \tag{8}$$

If the eigenspace decomposition $\sum_{s\in I}\oplus\mathcal{A}_s$ of $\mathrm{ad}(\mu^{-1})$ is given, then setting $\mathcal{L}_s{=}\mathcal{A}_s/L\cap\mathcal{A}_s$, we have

$$\mu^{-1}{*}\phi_s{*}\varpi=(s{+}c^{-1})\phi_s{*}\varpi, \quad \mu{*}\phi_s{*}\varpi=\frac{c}{cs{+}1}\phi_s{*}\varpi, \quad \phi_s\in\mathcal{L}_s.$$

As μ^{-1} is viewed as the energy, the positivity $s{+}c^{-1}{>}0$ may be required, but there is no effective relation between ϖ and ϕ_s.

The observation above suggests that the next target of the theory of deformation quantization is to construct μ-regulated algebras with abstract vacuums.

As $1{=}\varpi{+}(1{-}\varpi)$ and $\mathcal{A}{=}\mathcal{A}{*}\varpi\oplus\mathcal{A}{*}(1{-}\varpi)$, we see

$$\mathcal{A}{=}\mathbb{C}\varpi\oplus(1{-}\varpi){*}\mathcal{A}{*}\varpi\oplus\varpi{*}\mathcal{A}{*}(1{-}\varpi)\oplus(1{-}\varpi){*}\mathcal{A}(1{-}\varpi).$$

In what follows we assume there exist closed linear subspaces $\mathcal{L}\subset(1{-}\varpi){*}\mathcal{A}$ and $\mathcal{R}\subset\mathcal{A}{*}(1{-}\varpi)$ such that $\mathcal{L}\cap\mathcal{A}^{-\infty}{=}\{0\}$, $\mathcal{R}\cap\mathcal{A}^{-\infty}{=}\{0\}$ and

$$(1{-}\varpi){*}\mathcal{L}{*}\varpi{=}(1{-}\varpi){*}\mathcal{A}{*}\varpi, \quad \varpi{*}\mathcal{R}{*}(1{-}\varpi){=}\varpi{*}\mathcal{A}{*}(1{-}\varpi)$$

and the projections

$$\pi:\mathcal{L}{=}(1{-}\varpi){*}\mathcal{L}\to(1{-}\varpi){*}\mathcal{L}{*}\varpi, \quad \pi:\mathcal{R}{=}\mathcal{R}{*}(1{-}\varpi)\to\varpi{*}\mathcal{R}{*}(1{-}\varpi)$$

are linear isomorphic. We set

$$\tilde{\mathcal{L}}{=}\mathbb{C}{*}1\oplus(1{-}\varpi){*}\mathcal{L}, \quad \tilde{\mathcal{R}}{=}\mathbb{C}{*}1\oplus\mathcal{R}{*}(1{-}\varpi).$$

Note that the projections $\pi:\tilde{\mathcal{L}}\to\tilde{\mathcal{L}}{*}\varpi$, $\pi:\tilde{\mathcal{R}}\to\varpi{*}\tilde{\mathcal{R}}$ are linear isomorphisms. As every $a{\in}\mathcal{A}$ is written by

$$a{=}\varpi{*}a{*}\varpi{+}(1{-}\varpi){*}a{*}\varpi{+}\varpi{*}a{*}(1{-}\varpi){+}(1{-}\varpi){*}a{*}(1{-}\varpi), \quad \varpi{*}a{*}\varpi{=}\delta_a\varpi{\in}\mathbb{C}\varpi,$$

we see that $\delta_a\varpi{+}(1{-}\varpi){*}a{*}\varpi\in\tilde{\mathcal{L}}{*}\varpi$, and $\delta_a{+}(1{-}\varpi){*}a\in\tilde{\mathcal{L}}$.

Proposition 2.2 *Let $L=\{f\in\mathcal{A}; f{*}\varpi{=}0\}$. Then $L=\varpi{*}\mathcal{A}{*}(1{-}\varpi){\oplus}(1{-}\varpi){*}\mathcal{A}(1{-}\varpi)$ and $\mathcal{A}/L$ is linearly isomorphic to $\tilde{\mathcal{L}}$ and $\tilde{\mathcal{L}}{*}\varpi$. An element a of $\mathcal{L}$ is characterized by $a{=}(1{-}\varpi){*}a$.*

Similarly, let $R=\{f\in\mathcal{A}; \varpi{*}f{=}0\}$. Then $R=(1{-}\varpi){*}\mathcal{A}{\oplus}(1{-}\varpi){*}\mathcal{A}(1{-}\varpi)$ and $R\backslash\mathcal{A}$ is linearly isomorphic to $\tilde{\mathcal{R}}$ and $\varpi{*}\tilde{\mathcal{L}}$. An element b of $\mathcal{R}$ is characterized by $b{=}b{*}(1{-}\varpi)$.

The regular representation, which is called often a **vacuum representation** of $\mathcal{A}$ is defined for $a{\in}\mathcal{A}$ by

$$P(a):\tilde{\mathcal{L}}\to\tilde{\mathcal{L}},\quad P(a)(\phi){=}\pi^{-1}(a{*}\phi{*}\varpi).$$

Since $\pi^{-1}(a{*}\varpi){*}\varpi{=}a{*}\varpi$, it is clear that $P(a)P(b){=}P(a{*}b)$. In particular,

$$P(\varpi)(\phi){=}\pi^{-1}(\varpi{*}\phi{*}\varpi){=}\delta_{\phi},\quad \delta_{\phi}{\in}\mathbb{C},\quad \delta_{\varpi}{=}1 \tag{9}$$

is a projection operator of rank one. In particular, $\tilde{\mathcal{L}}{*}\tilde{\mathcal{L}}{*}\varpi{=}\tilde{\mathcal{L}}{*}\varpi$, but $\tilde{\mathcal{L}}{*}\tilde{\mathcal{L}}{=}\tilde{\mathcal{L}}$ does not hold in general.

On the other hand, there is a natural bilinear form $\langle\ ,\ \rangle_{\varpi}:R\backslash\mathcal{A}\times\mathcal{A}/L\to\mathbb{C}$ over $\mathbb{C}$ defined by

$$\varpi{*}\psi{*}\phi{*}\varpi{=}\langle\psi{*}\phi\rangle_{\varpi}\varpi. \tag{10}$$

This is identified with the bilinear form

$$\langle\ ,\ \rangle_{\varpi}:\tilde{\mathcal{R}}\times\tilde{\mathcal{L}}\to\mathbb{C}. \tag{11}$$

In this note, we mainly consider the matrix representation given in the form $\tilde{\mathcal{L}}{*}\varpi{*}\tilde{\mathcal{R}}$. Obviously, we have $\tilde{\mathcal{L}}{*}\varpi{*}\tilde{\mathcal{R}}\subset\mathcal{A}^{-\infty}$ and

$$(\tilde{\mathcal{L}}{*}\varpi{*}\tilde{\mathcal{R}}){*}(\tilde{\mathcal{L}}{*}\varpi{*}\tilde{\mathcal{R}}){=}\tilde{\mathcal{L}}{*}\varpi{*}\tilde{\mathcal{R}}.$$

In typical examples, the bilinear form (11) is non-degenerate, and there are linear basis

$$\tilde{\mathcal{L}}:e_0,e_1,e_2,\cdots,e_k,\cdots,\quad \tilde{\mathcal{R}}:f_0,f_1,f_2,\cdots,f_k,\cdots,\quad e_0{=}1{=}f_0,$$

such that $\langle e_i,f_j\rangle_{\varpi}{=}\delta_{ij}$. In this case, we see $e_k{*}\varpi{*}f_{\ell}$ is the (k,ℓ) matrix element and $\varpi{=}e_0{*}\varpi{*}f_0$. They are elements of $\mathcal{A}^{-\infty}$, but the representation space is spanned by $\{e_k{*}\varpi;k=0,1,2,\cdots\}$ together with certain topologies.

3 Extended Weyl algebra

Elements such as abstract vacuums are transcendental elements. For the systematic treatment for such elements, we have to begin with the transcendental extension of Weyl algebra.

Let $\boldsymbol{u}{=}(u_1,\cdots,u_m,u_{m+1},\cdots,u_{2m})$. For an arbitrary fixed $2m{\times}2m$-complex symmetric matrix $K{\in}\mathcal{S}_{\mathbb{C}}(2m)$, we set $\Lambda{=}K{+}J$ where J is the standard skew-symmetric matrix $J{=}\begin{bmatrix}0 & -I_m\\ I_m & 0\end{bmatrix}$. We define a product $*_{_K}$ on the space of polynomials $\mathbb{C}[\boldsymbol{u}]$ by the formula

$$f*_{_K}g=fe^{\frac{i\hbar}{2}(\sum\overleftarrow{\partial_{u_i}}\Lambda^{ij}\overrightarrow{\partial_{u_j}})}g=\sum_k\frac{(i\hbar)^k}{k!2^k}\Lambda^{i_1j_1}{\cdots}\Lambda^{i_kj_k}\partial_{u_{i_1}}{\cdots}\partial_{u_{i_k}}f\ \partial_{u_{j_1}}{\cdots}\partial_{u_{j_k}}g, \tag{12}$$

where $i\hbar\in\mathbb{C}\backslash\{0\}$ is a complex parameter. When $K{=}0$, the product formula (12) is called the **Moyal product formula**. (In [15], [16], [17], [18], [19], $*_{_K}$-product is first denoted by $*_{_\Lambda}$, or $*_{_{K+J}}$, and then the notation $*_{_K}$ is used after fixing the skew part J.)

It is known and not hard to prove that $(\mathbb{C}[\boldsymbol{u}],*_{_K})$ are mutually isomorphic associative algebras for every K. The Weyl algebra $(W_{2m}[\hbar];*)$ is the name of the isomorphism class.

In calculations in the algebra where we need only the commutation relations, we have no need to use the expression by K. Thus, it is natural to think that the algebraic essence must be free from their expressions, which may be called *the independence of ordering principle* (IOP in short). But this is a dangerous trip. In highly transcendental elements, the expression parameter K plays an essential role (cf.§, 3.1.1). We have seen in [16] that the expression parameter is essential even in the one variable commutative case.

If K is fixed, the $*_{_K}$-product formula gives a way of univalent expression for elements of $(W_{2m}[\hbar];*)$. Every element $f_*{\in}(W_{2m}[\hbar];*)$ is expressed in the form of ordinary polynomial, which we denote by $:\!f_*\!:_{_K}\in\mathbb{C}[\boldsymbol{u}]$ and call this **the K-ordered expression (or K-expression in brief) of f_***. In this context, the product formula (12) is often called a K-ordered expression. $K=(K^{ij}){=}(\langle u_i, Ku_j\rangle)$ is called an expression parameter. Via univalent expressions, one can consider topological completion of the algebra, and various transcendental elements.

The intertwiner between K-ordered expression and K'-ordered expression is explicitly given as follows:

Proposition 3.1 *For every $K, K'\in\mathcal{S}_{\mathbb{C}}(2m)$, the intertwiner is defined by*

$$I_{_K}^{^{K'}}(f)=\exp\Big(\frac{i\hbar}{4}\sum_{i,j}(K'^{ij}{-}K^{ij})\partial_{u_i}\partial_{u_j}\Big)f\ \ (=I_{_0}^{^{K'}}(I_{_0}^{^K})^{-1}(f)), \tag{13}$$

which gives an isomorphism $I_{_K}^{^{K'}}:(\mathbb{C}[\boldsymbol{u}];*_{_K})\to(\mathbb{C}[\boldsymbol{u}];*_{_{K'}})$. Namely, for any $f,g\in\mathbb{C}[\boldsymbol{u}]$, we have

$$I_{_K}^{^{K'}}(f*_{_K}g)=I_{_K}^{^{K'}}(f)*_{_{K'}}I_{_K}^{^{K'}}(g). \tag{14}$$

Intertwiners do not change the algebraic structure $*$, but these change the expression of elements by the ordinary commutative structure.

The next formula is trivial, but often very useful in the concrete calculation: (Cf.§ 3.3.1.) Suppose $K=K_1{+}K_2$. Then

$$I_{_{K_2}}^{^{K_1+K_2}}I_{_0}^{^{K_2}}=I_{_0}^{^{K_1+K_2}}. \tag{15}$$

Example 1 Let $Hol(\mathbb{C}^{2m})$ (resp. $C^\infty(\mathbb{R}^{2m})$) be the space of all holomorphic (resp. C^∞-) functions on $\mathbb{C}^{2m}$ (resp. $\mathbb{R}^{2m}$). By setting $\nu=\hbar$ in (12), and by fixing the expression parameter K in $\mathcal{S}_{\mathbb{C}}(2m)$, we see that the Heisenberg algebra $\mathcal{H}_{2m}[\nu]$ is a ν-regulated algebra.

Let $:\!\mathcal{A}\!:_{_K}=Hol(\mathbb{C}^{2m})[[\nu]]$, (resp, $:\!\mathcal{A}\!:_{_K}=C^\infty(\mathbb{R}^{2m})[[\nu]]$), the space of all formal power series of ν. Then, (12) makes $(:\!\mathcal{A}\!:_{_K};*_{_K})$ an associative algebra. As they are all mutually isomorphic by the intertwiners, the isomorphism class will be denoted by $(\mathcal{A};*)$. This is a ν-regulated algebra such that $[\nu,\mathcal{A}]{=}\{0\}$ and

$$\mathcal{A}{=}Hol(\mathbb{C}^{2m}){\oplus}\nu{*}\mathcal{A},\quad \textit{resp.}\quad \mathcal{A}{=}C^\infty(\mathbb{R}^{2m}){\oplus}\nu{*}\mathcal{A},$$

involving abstract symplectic structure. The isomorphism class $(\mathcal{A};*)$ will be called the **formally extended Weyl algebra**, or the **formally extended Heisenberg algebra**.

Although the proof will be skipped, the next one is an important remark:

Proposition 3.2 *Changing expression parameter K corresponds to the replacement of complementary subspace B. In fact, the latter is much wider than changing K.*

3.1 Extension of products

For every positive real number p, we set

$$\mathcal{E}_p(\mathbb{C}^{2m}) = \{f \in \mathit{Hol}(\mathbb{C}^{2m})\,;\, \|f\|_{p,s} = \sup |f|\, e^{-s|\boldsymbol{u}|^p} < \infty,\ \forall s > 0\} \tag{16}$$

where $\boldsymbol{u} = (u_1, \dots, u_{2m})$ and $|\boldsymbol{u}| = (\sum_i |u_i|^2)^{1/2}$. The family of seminorms $\{||\cdot||_{p,s}\}_{s>0}$ induces a topology on $\mathcal{E}_p(\mathbb{C}^{2m})$ and $(\mathcal{E}_p(\mathbb{C}^{2m}), \cdot)$ is an associative commutative Fréchet algebra, where the dot $\cdot$ is the ordinary product for functions in $\mathcal{E}_p(\mathbb{C}^{2m})$. It is easily seen that for $0 < p < p'$, there is a continuous embedding

$$\mathcal{E}_p(\mathbb{C}^{2m}) \subset \mathcal{E}_{p'}(\mathbb{C}^{2m}) \tag{17}$$

as commutative Fréchet algebras (cf. [6]), and that $\mathcal{E}_p(\mathbb{C}^{2m})$ is $GL(n, \mathbb{C})$-invariant.

We denote

$$\mathcal{E}_{p+}(\mathbb{C}^{2m}) = \bigcap_{p'>p} \mathcal{E}_{p'}(\mathbb{C}^{2m}), \quad \text{(with the intersection topology)} \tag{18}$$

It is obvious that every polynomial is contained in $\mathcal{E}_p(\mathbb{C}^{2m})$, that is $p(\boldsymbol{u}) \in \mathcal{E}_{0+}(\mathbb{C}^{2m})$, and $\mathbb{C}[\boldsymbol{u}]$ is dense in $\mathcal{E}_p(\mathbb{C}^{2m})$ for any $p > 0$ in the Fréchet topology defined by the family of seminorms $\{||\ ||_{p,s}\}_{s>0}$.

We easily see that $e^{\sum a_i u_i} \in \mathcal{E}_{1+}(\mathbb{C}^{2m})$. Moreover, it is not difficult to show that an exponential function $e^{p(\boldsymbol{u})}$ of a polynomial $p(\boldsymbol{u})$ of degree d is contained in $\mathcal{E}_{d+}(\mathbb{C}^{2m})$, but not in $\mathcal{E}_d(\mathbb{C}^{2m})$.

Theorem 3.1 (*Cf.*[13],[14]) *For $0 < p \leq 2$, the product formula* (12) *extends to give the following:*
(1) *The space $(\mathcal{E}_p(\mathbb{C}^{2m}), *_{K})$ forms a complete noncommutative topological associative algebra.*
(2) *The intertwiner $I_{K}^{K'}$ extends to give an isomorphism of $(\mathcal{E}_p(\mathbb{C}^{2m}), *_{K})$ onto $(\mathcal{E}_p(\mathbb{C}^{2m}), *_{K'})$.*

Hence we denote the isomorphism class by $(\mathcal{E}_p(\mathbb{C}^{2m}), *)$ $(0 \leq p \leq 2)$, and we call this the *extended Weyl algebra*. This is naturally an $\hbar$-regulated algebra. An element H_* of $(\mathcal{E}_p(\mathbb{C}^{2m}), *)$ is a family $\{H_{K}; K \in \mathcal{S}_{\mathbb{C}}(2m)\}$ such that $I_{K}^{K'} H_{K} {=} H_{K'}$, but it is convenient to denote H_{K} by $:H_*:_{K}$.

Note that the exponential function of a quadratic form can not be treated by Theorem 3.1.

In the case $p{>}2$, we see the next one:

Theorem 3.2 (*Cf.*[13],[14]) *For $p{>}2$, find p' such that $\frac{1}{p} + \frac{1}{p'} {\geq} 1$, then the product* (12)*extends to a continuous bilinear mapping*

$$\mathcal{E}_p(\mathbb{C}^{2m}) \times \mathcal{E}_{p'}(\mathbb{C}^{2m}) \to \mathcal{E}_p(\mathbb{C}^{2m}), \quad \mathcal{E}_{p'}(\mathbb{C}^{2m}) \times \mathcal{E}_p(\mathbb{C}^{2m}) \to \mathcal{E}_p(\mathbb{C}^{2m}). \tag{19}$$

If two of $f, g, h \in \mathcal{E}_p(\mathbb{C}^{2m})$ $(p{>}2)$ are in $\mathcal{E}_{p'}(\mathbb{C}^{2m})$, then the associativity $(f{*_{K}}g){*_{K}}h = f{*_{K}}(g{*_{K}}h)$ holds. Namely, $\mathcal{E}_p(\mathbb{C}^{2m})$ is a two-sided $\mathcal{E}_{p'}(\mathbb{C}^{2m})$-module.

3.1.1 Several remarks on the notation $:A_*:_{K}$

In previous notes we had used notations such as $:A_*:_{K}$, and it will be used often in what follows. Here we have to explain what A_* is. This is in fact the family $\{A_{K}; K \in \mathcal{I}\}$ where A_{K} are elements computed by K-expression which are mutually intertwined by $I_{K}^{K'}$ for every $K, K' \in \mathcal{I}$. $\mathcal{I}$ is a subset of $\mathcal{S}_{\mathbb{C}}(2m)$. If $\mathcal{I}$ contains an open dense subset of $\mathcal{S}_{\mathbb{C}}(2m)$, then we say that A_* is given by a generic ordered expression.

Sometimes, elements such as A_*^{-1} or $\sum_{k\in\mathbb{Z}} e_*^{kA}$, $\mathcal{I}$ is not a dense subset. It contains only a nonempty open subset. Even in such a case we use notations $:A_*:_{_K}$ by indicating the domain $\mathcal{I}$ for A_*.

It is sometimes very convenient to compute under a fixed expression parameter K. In particular, Weyl ordered ($K=0$) expression and the normal ordered ($K=\left[\begin{smallmatrix}0&1\\1&0\end{smallmatrix}\right]$) expression are very convenient to compute. However, such expressions are not generic. Therefore, the results might not be a generic result.

However, the next Proposition shows that such a restricted expression can be a useful tool.

Proposition 3.3 *Suppose we have an identity* $:C_*:_{_{K_0}}{=}:A_*:_{_{K_0}}*_{_{K_0}}:B_*:_{_{K_0}}$ *under a fixed* K_0*, and suppose* $:A_*:_{_{K_t}}$*,* $:B_*:_{_{K_t}}$ *and their* $*_{_{K_t}}$ *product* $:A_*:_{_{K_t}}*_{_{K_t}}:B_*:_{_{K_t}}$ *are defined on a path* K_t*,* $t{\in}[0,1]$*. Then, one may define* $:C_*:_{_{K_1}}$ *by* $:C_*:_{_{K_1}}{=}:A_*:_{_{K_1}}*_{_{K_1}}:B_*:_{_{K_1}}$*.*

In general, the r.h.s. depends on the path to K_0 to K_1. Indeed, it causes the double valued nature of the $*$-exponential functions of quadratic forms. Thus, a transcendental element A_* in a $*$-algebra is defined always with expression parameters $\mathcal{I}(A_*)$, and sometimes with the path from the reference point (cf. the notion of synchronized path selection in [18]).

If $\mathcal{I}(A_*)$ contains open dense subset, then it may not be explicitly indicated. However, $\mathcal{I}(A_*)$ might be restricted in a very small region. Furthermore, in [17], [19], we saw that a certain family of elements behaves very different way under some special expression parameters.

3.2 Star-exponential functions

The $*$-exponential function $e_*^{tH_*}$ for $H_*{\in}(\mathcal{A},*)$ is defined as the collection $:e_*^{tH_*}:_{_K}$ of its K-expressions. If $\hbar$ is a formal parameter, then $:e_*^{tH_*}:_{_K}$ is defined as a series $\sum\frac{t^n}{n!}:H_*^n:_{_K}$ formally rearranged as power series of $\hbar$. In the case $\hbar$ is not formal, the $*$-exponential function $e_*^{tH_*}$ is not defined by a power series. If H_* is a polynomial of degree 3, then the radius of convergence is 0 in general (cf. [16]).

Instead, these are considered as the solution of an evolution equation

$$\frac{d}{dt}:f_t:_{_K}{=}:H_*:_{_K}*_{_K}:f_t:_{_K}, \quad :f_0:_{_K}{=}1.$$

This is a differential equation, if H_* is a polynomial, but the solution may not exist in general, but a real analytic solution is unique. The solution (if exists uniquely) with initial data $:g_*:_{_K}$ will be denoted by $:e_*^{tH_*}{*}g_*:_{_K}$.

Note however even though $:e_*^{tH_*}$ is not defined by itself, there is a case that $:e_*^{tH_*}{*}g_*:_{_K}$ is welldefined for some initial data. Moreover, even though $:e_*^{tH_*}{*}g_*:_{_K}$ is not defined, there is a case where $:e_*^{tH_*}{*}g_*{*}e_*^{-tH_*}:_{_K}$ is welldefined as $e^{t\mathrm{ad}(H_*)}g_*$.

To simplify notations, we often denote $\langle \boldsymbol{a}\Gamma,\boldsymbol{b}\rangle=\sum_{ij=1}^{2m}\Gamma^{ij}a_ib_j$, $\langle\boldsymbol{a},\boldsymbol{u}\rangle=\sum_{i=1}^{2m}a_iu^i$. These will be denoted also by $\boldsymbol{a}\Gamma\,{}^t\boldsymbol{b}$ and $\langle\boldsymbol{a},\boldsymbol{u}\rangle=\boldsymbol{a}\,{}^t\boldsymbol{u}$. It is easy to see that

$$[\langle\boldsymbol{a},\boldsymbol{u}\rangle,\langle\boldsymbol{b},\boldsymbol{u}\rangle]_*=i\hbar\langle\boldsymbol{a}J,\boldsymbol{b}\rangle.$$

If H_* is a linear function, then this is given also as follows: By a direct calculation of intertwiner, we see that

$$I_{_K}^{^{K'}}(e^{\frac{1}{i\hbar}\langle\boldsymbol{a},\boldsymbol{u}\rangle})=e^{\frac{1}{4i\hbar}\boldsymbol{a}(K'-K)\,{}^t\boldsymbol{a}}e^{\frac{1}{i\hbar}\langle\boldsymbol{a},\boldsymbol{u}\rangle}. \tag{20}$$

Hence, $\{e^{\frac{1}{4i\hbar}\boldsymbol{a}K\,{}^t\!\boldsymbol{a}}e^{\frac{1}{i\hbar}\langle\boldsymbol{a},\boldsymbol{u}\rangle}; K\in\mathcal{S}_{\mathbb{C}}(2m)\}$ is a family of mutually isomorphic one parameter groups cf. [16],[17],[18]. We have denoted this element symbolically by $e_*^{\frac{1}{i\hbar}\langle\boldsymbol{a},\boldsymbol{u}\rangle}$. Namely we denote

$$:e_*^{\frac{1}{i\hbar}\langle\boldsymbol{a},\boldsymbol{u}\rangle}:_{K}= e^{\frac{1}{4i\hbar}\langle\boldsymbol{a}K,\boldsymbol{a}\rangle}e^{\frac{1}{i\hbar}\langle\boldsymbol{a},\boldsymbol{u}\rangle} = e^{\frac{1}{4i\hbar}\langle\boldsymbol{a}K,\boldsymbol{a}\rangle+\frac{1}{i\hbar}\langle\boldsymbol{a},\boldsymbol{u}\rangle}. \tag{21}$$

The fact stated below is an example how a transcendental element depends on the expression parameters:

Proposition 3.4 *If* $\mathrm{Im}\langle\boldsymbol{a}K,\boldsymbol{a}\rangle$ *is negative, then the formula of Fourier transform gives*

$$\int_{-\infty}^{\infty}:e_*^{t\frac{1}{i\hbar}\langle\boldsymbol{a},\boldsymbol{u}\rangle}:_{K}\,dt\in\mathcal{E}_{2+}(\mathbb{C}^{2m}).$$

In particular, both

$$:\langle\boldsymbol{a},\boldsymbol{u}\rangle:_{K+}^{-1}=\frac{1}{i\hbar}\int_{-\infty}^{0}:e_*^{t\frac{1}{i\hbar}\langle\boldsymbol{a},\boldsymbol{u}\rangle}:_{K}\,dt,\quad :\langle\boldsymbol{a},\boldsymbol{u}\rangle:_{K-}^{-1}=-\frac{1}{i\hbar}\int_{0}^{\infty}:e_*^{t\frac{1}{i\hbar}\langle\boldsymbol{a},\boldsymbol{u}\rangle}:_{K}\,dt$$

are $*$*-inverses of* $:\langle\boldsymbol{a},\boldsymbol{u}\rangle:_{K}$ *under the calculation of* $*_{K}$*-product.* $:\langle\boldsymbol{a},\boldsymbol{u}\rangle:_{K\pm}^{-1}$ *belongs to* $\mathcal{E}_{2+}(\mathbb{C}^{2m})$*. Two different inverses of a single element breaks apparently the associativity.*

Furthermore, if $\mathrm{Im}\langle\boldsymbol{a}K,\boldsymbol{a}\rangle$ is negative, then $\sum_{n=-\infty}^{\infty}:e_*^{n\frac{1}{i\hbar}\langle\boldsymbol{a},\boldsymbol{u}\rangle}:_{K}$ converges to give Jacobi's theta function $\Theta_3(\frac{1}{i\hbar}\langle\boldsymbol{a},\boldsymbol{u}\rangle)$ in [16].

Anti-automorphism and Hermite structure

Note that generators of the algebra $(W_{2m},*)$ are only abstract symbols. It does not make sense to say that these are complex variables or not. To make it clear we fix an hermitian structure by introducing an anti-automorphism $x\to x^{\iota}$, called the **Hermitian conjugate**, by defining

$$u_k^{\iota}{=}u_k,\quad k=1{\sim}2m,\quad i^{\iota}=-i,\quad \hbar^{\iota}=\hbar.$$

An element $H\in W_{2m}$ is called an *hermite element* when $H^{\iota}{=}H$, but its K-ordered expression $:H_*:_{K}$ does not have the property $\overline{:H_*:_{K}}{=}:H_*:_{K}$ in general.

The next identity is useful.

Proposition 3.5 *If* $e_*^{tH_*}$, $e_*^{tH_*^{\iota}}$ *and* $e_*^{sH_*^{\iota}}{*}e_*^{tH_*}$ *are defined for* $s,t\in\mathbb{R}$*, then* $e_*^{tH_*^{\iota}}{=}(e_*^{tH_*})^{\iota}$*. In particular,* $e_*^{tu_i}$ *is an hermite element for every* $t\in\mathbb{R}$*.*

Proof. Set $f_t=e_*^{tH_*}$, $g_t=e_*^{tH_*^{\iota}}$. Then $\frac{d}{dt}f_{-t}{=}(-H_*){*}f_{-t}$. It follows $\frac{d}{dt}f^{\iota}_{-t}{=}f^{\iota}_{-t}{*}(-H^{\iota}_*)$. Thus, we have

$$\frac{d}{dt}f^{\iota}_{-t}{*}g_t{=}f^{\iota}_{-t}{*}(-H^{\iota}_*{+}H^{\iota}_*){*}g_t{=}0.$$

It follows $e_*^{tH_*^{\iota}}{=}(e_*^{tH_*})^{\iota}$ as $f^{\iota}_{-t}{*}g_t{=}1$. □

If H_* is hermitian and $F_t{=}e_*^{itH_*}$ exists, then F_t is unitary, since we have the primitive conservation law:

$$\frac{d}{dt}{F^{\iota}}_t{*}F_t=0,\quad {F^{\iota}}_0{*}F_0=1. \tag{22}$$

3.2.1 The case $\rho \in \mathcal{H}_{2m}[\nu]$

Suppose first that ρ mentioned in § 1 satisfies $\rho{\in}\mathcal{H}_{2m}[\nu]$, then ρ must be a linear function

$$\langle \boldsymbol{a}, \tilde{\boldsymbol{u}}\rangle{=}\sum_{i=1}^{m}(a_i u_i{+}a_{m+i}v_i).$$

By Proposition 3.4, if $\mathrm{Im}\langle \boldsymbol{a}K, \boldsymbol{a}\rangle$ is negative, then $\langle \boldsymbol{a}, \tilde{\boldsymbol{u}}\rangle$ is invertible.

Suppose $\rho{=}u_1$ for simplicity and denote its inverse by $u_{1\circ}^{-1}$. Then, the algebra generated by (3) in § 1 turns out to be generated by

$$u_{1\circ}^{-2}{*}\nu,\ u_{1\circ}^{-1}{*}v_1,\ u_{1\circ}^{-1}{*}u_2,\ \cdots,\ u_{1\circ}^{-1}{*}u_m,\ u_{1\circ}^{-1}{*}v_2,\ \cdots,\ u_{1\circ}^{-1}{*}v_m \tag{23}$$

where the calculations are executed by the $*_{_K}$-product formula under the K-ordered expression. Thus, the suffix such as $*_{_K}$, $:\ :_{_K}$ are omitted.

For simplicity we denote these as follows:

$$\mu,\ \tau,\ \tilde{u}_k,\ \tilde{v}_k,\ k{=}2{\sim}m$$

and let $\mathcal{A}$ be the algebra generated by these in $\mathcal{E}_{2+}(\mathbb{C}^{2m})$. Using $[u_{1\circ}^{-1}, v_1]{=}i\hbar u_{1\circ}^{-2}$, $[u_{1\circ}^{-1}, u_k]{=}0{=}[u_{1\circ}^{-1}, v_k]$, $k = 2{\sim}m$, we have

$$[\mu^{-1}, \tau]{=}2i, \quad [\tilde{u}_i,\ \tilde{v}_j]{=}\mu i\delta_{ij}, \quad [\tau, \tilde{u}_i]{=}\mu i\tilde{u}_i, \quad [\tau, \tilde{v}_i]{=}\mu i\tilde{v}_i, \quad \text{all others are 0.}$$

As $[\mu, \tilde{u}_i]{=}0{=}[\mu, \tilde{v}_i]$, $i{=}2{\sim}m$, $\{\mu, \tilde{u}_i, \tilde{v}_i\}$ generates the Heisenberg algebra $\mathcal{H}_{2m}[\mu]$. Hence $\mathcal{A}$ is a μ-regulated algebra $\mathcal{H}_{2m}[\mu, \tau]$. As $[\tau, \mu^{-1}]{=}-2i$, τ may be viewed as a canonical conjugate of μ^{-1}.

Now join u_1 to $\mathcal{H}_{2m}[\mu, \tau]$. This is also a μ-regulated algebra.

Proposition 3.6 *A suitably extended μ-regulated algebra such as $\mathcal{H}_{2m}[\mu, u_1, \tau]$ contains no standard abstract vacuum* (*cf.* § *2.1*).

Proof. Suppose there is a standard abstract vacuum ϖ in $\varpi \in \mathcal{H}_{2m}[\mu, u_1, \tau]$. It follows $\varpi{*}u_1{=}0$ or $u_1{*}\varpi{=}0$. Hence we have $\frac{d}{ds}\varpi{*}e_*^{su_1}{*}\varpi{=}0$ and hence $\varpi{*}e_*^{su_1}{*}\varpi{=}\varpi$. Now recall the formula

$$u_{1\circ}^{-1}{*}u_{1\circ}^{-1}{=}\int_{-\infty}^{0} 2s{:}e_*^{su_1}{:}_{_K}ds.$$

Using this we have

$$\varpi{*}u_{1\circ}^{-1}{*}u_{1\circ}^{-1}{*}\varpi{=}\varpi *\int_{-\infty}^{0} 2se_*^{su_1}{*}\varpi ds{=}\int_{-\infty}^{0} 2s\varpi{*}e_*^{su_1}{*}\varpi ds{=}\infty\varpi.$$

It follows $\varpi{*}\mu{*}\varpi{=}\infty\varpi$. □

Apparently, this is caused by the choice $u_{1\circ}^{-1}{\in}\mathcal{E}_{2+}(\mathbb{C}^{2m})$. If this is replaced by the embedded Heisenberg algebra given in § 3.2.2, then we can find an abstract vacuum in it (see § 4).

3.2.2 Heisenberg algebra embedded in Weyl algebra

In what follows we denote by

$$x_0, x_1, \cdots, x_m, y_0, y_1, \cdots, y_m \tag{24}$$

natural coordinate functions of $\mathbb{C}^{2m+2}$. Consider the Weyl algebra $(W_{2m+2}[\hbar], *)$ generated by setting the relation $[x_i, y_j] = -i\hbar\delta_{ij}$. Fix a generic expression parameter $K \in \mathcal{S}_{\mathbb{C}}(2m{+}2)$. In what follows, all calculations executes under the K-ordered expression and $*_{_K}$-product formula. We first embed the Heisenberg algebra $\mathcal{H}_{2m}[\nu]$ into $(\mathcal{E}_{1+}(\mathbb{C}^{2m+2}, *_{_K})$.

In this algebra, we make first the $*$-exponential function $e_*^{ty_0}$. As it was seen in § 3.2, the exponential law $e_*^{sy_0}{*}e_*^{ty_0}{=}e_*^{(s+t)y_0}$ holds and the K-ordered expression is $:e_*^{ty_0}:_{_K}{=}e^{\frac{i\hbar}{4}\delta'}e^{e^{ty_0}}$ by using a suitable constant δ'. Hence $e_*^{sy_0}$ is an element of $\mathcal{E}_{1+}(\mathbb{C}^{2m+2})$.

Set $\mu = \hbar{*}e_*^{-2y_0}$. As $\hbar{>}0$, the exponential law and Theorem 3.1 show that $\hbar e_*^{-2y_0}{*} : \mathcal{E}_{1+}(\mathbb{C}^{2m+2}) \to \mathcal{E}_{1+}(\mathbb{C}^{2m+2})$ is a linear isomorphism.

We set furthermore

$$\mu = \hbar e_*^{-2y_0}, \quad \tau = \frac{1}{2}(e_*^{-2y_0}{*}x_0{+}x_0{*}e_*^{-2y_0}), \quad \tilde{u}_i = e_*^{-y_0}{*}x_i, \quad \tilde{v}_i = e_*^{-y_0}{*}y_i, \quad i = 1 \sim m. \tag{25}$$

They are all hermite elements (cf.§ 3.2) and

$$\tau = e_*^{-2y_0}{*}(x_0{+}i\hbar){=}(x_0{-}i\hbar){*}e_*^{-2y_0},$$

as $[e_*^{-2y_0}, x_0] = -2i\hbar e_*^{-2y_0}{=}-2i\mu$. Using these we have

$$[\tau, \mu] = 2i\mu_*^2, \quad [\mu, \tilde{u}_i]{=}0, \quad [\mu, \tilde{v}_i]{=}0, \quad [\tau, \tilde{u}_i] = \mu i{*}\tilde{u}_i, \quad [\tau, \tilde{v}_i] = \mu i{*}\tilde{v}_i, \quad [\tilde{u}_i, \tilde{v}_j] = -i\mu\delta_{ij}. \tag{26}$$

As $[\mu{*}\mu^{-1}, \tau] = 0$, we see $[\mu^{-1}, \tau] = 2i$. The algebra generated by these will be denoted by $\widetilde{\mathcal{H}}_{2m}[\mu, \tau]$. It is clear that $\widetilde{\mathcal{H}}_{2m}[\mu, \tau]$ contains the Heisenberg algebra. Since $\mu^{-1}{\circ}\tau = \frac{1}{\hbar}x_1$, E_t is defined as the automorphism $\mathrm{Ad}(e_*^{t\frac{1}{i\hbar}x_1})$.

On the other hand, note that $[\hbar^{-1}e_*^{2y_0}, e_*^{-2y_0}{*}x_0]{=}2i$, although $e_*^{-2y_0}{*}x_0$ is not hermitian. Hence $\hat{\tau}{=}e_*^{-2y_0}{*}x_0$ may be regarded as a canonical conjugate of μ^{-1}. Since $\mu^{-1}{*}\hat{\tau} = \frac{1}{\hbar}x_0$, we see $\mathrm{Ad}(e_*^{t\frac{1}{i\hbar}x_0}){=}E_t$. Thus, by using $\hat{\tau}$ instead of τ, we have the same commutation relations as (26).

Summarizing these we see the next

Theorem 3.3 *Heisenberg algebra $\mathcal{H}_{2m}[\mu]$ is isomorphic to the subalgebra of $\widetilde{\mathcal{H}}_{2m}[\mu, \tau]$ $(\subset \mathcal{E}_{1+}(\mathbb{C}^{2m+2}))$ where μ corresponds to $\hbar e_*^{-2y_0}$, and $\frac{1}{2}(e_*^{-2y_0}{*}x_0{+}x_0{*}e_*^{-2y_0})$ is a canonical conjugate of μ_*^{-1}.*

On the other hand, $\mathcal{H}_{2m}[\mu]$ is also isomorphic to the subalgebra of $\widetilde{\mathcal{H}}_{2m}[\mu, \hat{\tau}]$, although $\hat{\tau}$ is not an Hermite element.

Besides the topological completeness, $\widetilde{\mathcal{H}}_{2m}[\mu, \tau]$ (resp. $\widetilde{\mathcal{H}}_{2m}[\mu, \hat{\tau}]$) satisfies (A.1)∼(A.4) *and all properties of contact Weyl algebra, where $\frac{1}{2}\partial_\tau$ (resp.$\frac{1}{2}\partial_{\hat{\tau}}$) is the characteristic vector field and $B{=}\mathcal{H}_{2m}[\tau]$ (resp.$\mathcal{H}_{2m}[\hat{\tau}]$),*

In § 4, a vacuum representation of $\widetilde{\mathcal{H}}_{2m}[\mu, \tau]$ will be given. It is rather surprising that $:e_*^{it\tau}:_0$ is defined by a difference-differential equation, and $\mathrm{Ad}(e_*^{it\tau})$ is defined only for $t{\geq}0$. Such a semi-group property is familiar in heat equation. But, this appears clearer in the calculation of $e_*^{it\tau}$ under the Moyal product formula.

3.3 $*$-exponentials of quadratic forms

To define $\sqrt[*]{\rho_*^2}$ mentioned in § 1, we have to use the formula of Laplace transform

$$\sqrt[*]{\rho_*^2}^{-1}=\frac{1}{\sqrt{\pi}}\int_0^\infty \frac{1}{\sqrt{t}}e_*^{-t\rho_*^2}dt.$$

Thus, we have to know first the formula of $*$-exponential functions of quadratic forms.

As $(W_{2m+2}[\hbar],*) = (W_2[\hbar],*)\otimes(W_{2m}[\hbar],*)$, main properties of Weyl algebra are explained in $(W_2[\hbar],*)$. Let $(W_2[\hbar],*)$ be the Weyl algebra generated by x, y.

Let $K=\begin{bmatrix}\delta & c\\ c & \delta'\end{bmatrix}$ be the expression parameter. In this section, we summarize properties of some of $*$-exponential functions of quadratic forms. The K-ordered expression of the $*$-exponential function $e_*^{t\frac{1}{i\hbar}2x\circ y}$ is given as follows: By setting $\Delta{=}e^t{+}e^{-t}{-}c(e^t{-}e^{-t})$, it is given by

$$:e_*^{t\frac{1}{i\hbar}2x\circ y}:_K=\frac{2}{\sqrt{\Delta^2-(e^t-e^{-t})^2\delta\delta'}}\;e^{\frac{1}{i\hbar}\frac{e^t-e^{-t}}{\Delta^2-(e^t-e^{-t})^2\delta\delta'}\Big((e^t-e^{-t})(\delta' x^2+\delta y^2)+2\Delta xy\Big)}. \tag{27}$$

Setting $\delta\delta' = \rho^2$, we see

$$\sqrt{\Delta^2-(e^t-e^{-t})^2\delta\delta'} = e^{-t}\sqrt{((1-c+\rho)e^{2t}+(1+c-\rho))((1-c-\rho)e^{2t}+(1+c+\rho))}. \tag{28}$$

Set $H_*{=}\frac{1}{i\hbar}2x{\circ}y$. By (27), we easily see the following under generic expressions:

(**a**) $e_*^{tH_*}$ is rapidly decreasing on $\mathbb{R}$ of $e^{-|t|}$ order.

(**b**) $e_*^{zH_*}$ is $4\pi i$-periodic $:e_*^{(z+4\pi i)H_*}:_K = :e_*^{zH_*}:_K$. In precise for generic K-expressions there are a, b depending on K such that $-\infty{<}a{<}b{<}\infty$ and

  (**b,1**) if $a < s < b$, then $:e_*^{(s+it)H_*}:_K$ is 2π-periodic.

  (**b,2**) if $s{<}a$ or $b{<}s$, then $:e_*^{(s+it)H_*}:_K$ is alternating 2π-periodic.

(**c**) $\lim_{t\to-\infty} :e^{-t}e_*^{tH_*}:_K$ is a nontrivial element denoted by $:\varpi_{00}:_K$. We call this the **vacuum**.

In precise, the path tending to $-\infty$ should be in the same sheet as in the origin 1, as $*$-exponential functions have double branched singularities as it is seen in (27). By the exponential law $e^te_*^{tH_*}{=}e_*^{t(H_*+1)}$, ϖ_{00} is idempotent, i.e. $\varpi_{00}{*}\varpi_{00}{=}\varpi_{00}$ and the bumping identity $y{*}(x{\circ}y){=}(x{\circ}y{+}i\hbar){*}y$ give the property that

$$y{*}\varpi_{00}{=}0{=}\varpi_{00}{*}x.$$

Similarly, a nontrivial element $\overline{\varpi} = \lim_{t\to\infty} e^te_*^{tH_*}$ is called the **bar-vacuum**. This is an idempotent element such that $x{*}\overline{\varpi}_{00}{=}0{=}\overline{\varpi}_{00}{*}y$.

(**d**) Let $w{=}x{+}iy$, $\overline{w}{=}x{-}iy$. Then $w{\circ}\overline{w}{=}x_*^2{+}y_*^2$, $[w,\overline{w}]{=}-2\hbar$. By a similar calculation gives (cf. [18])

$$:e_*^{t\frac{1}{\hbar}(x_*^2+y_*^2)}:_K=\frac{1}{\sqrt{\Delta_K(t)}}e^{\frac{1}{i\hbar}\frac{i\sinh t}{\Delta_K(t)}\Big((\cosh t-\delta' i\sinh t)u^2+(\cosh t-\delta i\sinh t)v^2+2ci(\sinh z)uv\Big)} \tag{29}$$

where $\Delta_K(t)=\cosh^2 t-(\delta+\delta')i\sinh t\cosh t+(c^2-\delta\delta')\sinh^2 t$.

In generic ordered expression the **complex vacuum** $:\varpi_{\mathbb{C}}:_K$ is defined by

$$:\varpi_{\mathbb{C}}:_K=\lim_{t\to-\infty} :e^{-t}e_*^{t\frac{1}{\hbar}w\circ\overline{w}}:_K. \tag{30}$$

This exists by (29) and this is an idempotent element satisfying $\overline{w}*\varpi_{\mathbb{C}}=0=\varpi_{\mathbb{C}}*w$, providing the path tending to $-\infty$ avoiding singular points belongs to the same sheet as in the origin 1.

In spite of the double valued nature caused by $\sqrt{\ }$, the exponential law holds by computations such as $\sqrt{\alpha}\sqrt{\beta}=\sqrt{\alpha\beta}$. If $K=0$, then it is called the **Weyl ordered expression**, and if $c=1, \delta=0=\delta'$, then it is called the **normal ordered expression**. These are often used in physics, but general K-expressions are not used in physics.

For a degenerate quadratic form, the formula of the $*$-exponential functions are given by the case of one variable, and hence

$$:e_*^{t\frac{1}{i\hbar}x_*^2}:_K=\frac{\sqrt{i\hbar}}{\sqrt{i\hbar-t\delta}}e^{\frac{i\hbar t}{i\hbar-t\delta}x^2}.$$

Similar to vacuums, there exists the limit

$$\lim_{t\to\infty} :\sqrt{t}e_*^{t\frac{1}{i\hbar}x_*^2}:_K=\frac{\sqrt{i\hbar}}{\sqrt{-\delta}}e^{-\frac{i\hbar}{\delta}x^2}$$

and $:e_*^{t\frac{1}{i\hbar}x_*^2}:_K*_K\frac{\sqrt{i\hbar}}{\sqrt{-\delta}}e^{-\frac{i\hbar}{\delta}x^2}=\frac{\sqrt{i\hbar}}{\sqrt{-\delta}}e^{-\frac{i\hbar}{\delta}x^2}$. However, as $\sqrt{t}e_*^{t\frac{1}{i\hbar}x_*^2}$ is not a one parameter subgroup, the limit is not idempotent, but $\lim_{t\to\infty}(\sqrt{t}e_*^{t\frac{1}{i\hbar}x_*^2})_*^2$ diverges.

3.3.1 Abstract vacuums on several variables

Let $x_0, x_1, \cdots, x_m, y_0, y_1, \cdots, y_m$ be generators of $W_{2m+2}[\hbar]$. We want to apply the formula for the case $m=1$ mentioned above to the $*$-exponential functions of quadratic forms of $2m$-variables.

Obviously, the K-expression $:f_*(x_i,y_i)*g(x_j,y_j):_K$ of $f_*(x_i,y_i)*g(x_j,y_j)$ is computed by using only submatrices relating (i,j) components. For a given K, let

$$K_{(i)}=\begin{bmatrix} K_{i,i} & K_{i,m+i}\\ K_{m+i,i} & K_{m+1,m+i}\end{bmatrix}, \quad K_{(i,j)}=\begin{bmatrix} K_{i,j} & K_{i,m+j}\\ K_{m+j,i} & K_{m+i,m+j}\end{bmatrix}.$$

Then it is clear that

$$:f_*(x_i,y_i)*g_*(x_j,y_j):_K=:f_*(x_i,y_i):_{K_{(i)}}*_{K_{(i,j)}}:g_*(x_j,y_j):_{K_{(j)}}.$$

Recall now how the product $e_*^{H_*}*F$ is defined. This is defined by the value at $t=1$ of the real analytic solution of

$$\frac{d}{dt}f_t=H_**f_t, \quad f_0=F.$$

$e_*^{H_{*k}}*\cdots*e_*^{H_{*2}}*e_*^{H_{*1}}*F$ are defined similarly. Such products may be called the *path connecting products*. Hence in general $e_*^{H_{*2}}*e_*^{H_{*1}}*1$ might be different from $e_*^{H_{*1}}*e_*^{H_{*2}}*1$, even if $[H_{*1},H_{*2}]_*=0$. Recalling § 3.1.1, the next Proposition is not trivial.

Proposition 3.7 *In generic ordered expression, we have*

$$e_*^{ta_1x_1\circ y_1}{*}e_*^{ta_2x_2\circ y_2}{*}\cdots{*}e_*^{ta_mx_m\circ y_m}{=}e_*^{t(a_1x_1\circ y_1+a_2x_2\circ y_2+\cdots+a_mx_m\circ y_m)}$$

if each component has no singular point on the interval $[0,t]$.

Recall the remark given in § 3.3.1 about how to apply the formulas in the case $m=1$ to the case $m\geq 2$.

In what follows, several abstract vacuums which can be joined to the algebra $\mathcal{E}_{1+}(\mathbb{C}^{2m+2})$ without trouble, although they are in $\mathcal{E}_{2+}(\mathbb{C}^{2m+2})$.

Let

$$\widetilde{\varpi}(L_0){=}\varpi_{00}{*}\widetilde{\varpi}(L){=}\varpi_{00}{*}\prod_{k=1}^{m}{*}\varpi_{00}(k),\quad \widetilde{\varpi}(\overline{L}_0){=}\overline{\varpi}_{00}{*}\widetilde{\varpi}(L){=}\overline{\varpi}_{00}{*}\prod_{k=1}^{m}{*}\varpi_{00}(k), \tag{31}$$

where $\varpi_{00}(k){=}\lim_{t\to-\infty}e_*^{t\frac{1}{i\hbar}x_k\circ y_k}$, $\quad\overline{\varpi}_{00}{=}\lim_{t\to\infty}e^{t\frac{1}{2}}e_*^{t\frac{1}{i\hbar}x_0\circ y_0}$. These have properties as follows:

$$y_k{*}\widetilde{\varpi}(L_0){=}0,\ (k{=}0{\sim}m),\quad x_0{*}\widetilde{\varpi}(\overline{L}_0){=}0{=}\widetilde{\varpi}(\overline{L}_0){*}y_0,\ y_k{*}\widetilde{\varpi}(\overline{L}_0){=}0,\ (k{=}1{\sim}m).$$

Similarly, on $\mathbb{C}^m$, the **complex vacuum** $\widetilde{\varpi}_{\mathbb{C}}$ is given by

$$\widetilde{\varpi}_{\mathbb{C}}{=}\lim_{s\to-\infty}e_*^{s\sum\zeta_i*\overline{\zeta}_i}. \tag{32}$$

This satisfies $\overline{\zeta}_i{*}\widetilde{\varpi}_{\mathbb{C}}{=}0$.

As it was mentioned in § 1, there are various abstract vacuums. As $\widetilde{\varpi}(\overline{L}_0){*}e_*^{y_0}{*}\widetilde{\varpi}(\overline{L}_0){=}\widetilde{\varpi}(\overline{L}_0)$, we see $e_*^{y_0}{*}\widetilde{\varpi}(\overline{L}_0)$ is an abstract vacuum such that by using $(x_0{+}i\hbar)e_*^{y_0}{=}e_*^{y_0}{*}x_0$

$$(x_0{+}i\hbar)e_*^{y_0}{*}\widetilde{\varpi}(\overline{L}_0){=}0{=}\tau{*}e_*^{y_0}{*}\widetilde{\varpi}(\overline{L}_0).$$

We denote this by

$$\widetilde{\varpi}_{y_0}(\overline{L}_0){=}e_*^{y_0}{*}\widetilde{\varpi}(\overline{L}_0) \tag{33}$$

In what follows, we use the same notations as in (25). Under the notation (25), $e_*^{is\tau}{*}\varpi_{00}$ has the property as follows:

Proposition 3.8 *For* $|2s\hbar|{<}1$, $\varpi_{00}{*}e_*^{-is\tau}{*}\varpi_{00}{=}\frac{1}{\sqrt{1{+}2s\hbar}}\varpi_{00}$. *In particular,* $\sqrt{1{+}2s\hbar}\,e_*^{is\tau}{*}\varpi_{00}$ *is an idempotent element such that*

$$\varpi_{00}{*}x_0{*}\sqrt{1{+}2s\hbar}e_*^{is\tau}{*}\varpi_{00}{=}0,\quad \varpi_{00}{*}y_0{*}\sqrt{1{+}2s\hbar}e_*^{is\tau}{*}\varpi_{00}{=}i\hbar\sqrt{1{+}2s\hbar}e_*^{is\tau}{*}\varpi_{00}.$$

In particular, $e_*^{-is\tau}{*}\widetilde{\varpi}(L_0)$ *is an abstract vacuum for* $|2s\hbar|{<}1$.

Proof Note that $\varpi_{00}{*}x_0=0$, $e_*^{-2y_0}{*}\varpi_{00}{=}\varpi_{00}$. Computing $\varpi_{00}{*}(\tau_*^n){*}\varpi_{00}$ gives

$$\varpi_{00}{*}(-is\tau_*)^n{*}\varpi_{00}{=}(\frac{1}{2})_n(-s\hbar)^n{*}\varpi_{00},\quad (a)_n=a(a{+}1)\cdots(a{+}n{-}1).$$

Hence

$$\sum_n\varpi_{00}{*}\frac{1}{n!}(-is\tau_*)^n{*}\varpi_{00}{=}\sum_n\frac{1}{n!}(\frac{1}{2})_n(-s\hbar)^n\varpi_{00}{=}\frac{1}{\sqrt{1{+}2s\hbar}}{*}\varpi_{00},\quad |2s\hbar|<1.$$

In (54) later, the condition $|2s\hbar|{<}1$ is relaxed to $1{+}2s\hbar>0$. □

We denote this by

$$\widetilde{\varpi}_s(\mathcal{H}){=}e_*^{-is\tau}{*}\widetilde{\varpi}(L_0). \tag{34}$$

$\widetilde{\varpi}_{y_0}(\overline{L}_0)$ and $\widetilde{\varpi}_s(\mathcal{H})$ will be used in § 4.

3.4 Reductions vs Restriction to the Energy surface

Consider the most familiar case $\rho_*^2{=}\sum_{k=1}^{m}(u_k{*}u_k{+}v_k{*}v_k)$ mentioned in § 1. Set

$$\zeta_k{=}u_k{+}iv_k, \quad \bar\zeta_k{=}u_k{-}iv_k, \quad k = 1{\sim}m.$$

Then, $\rho_*^2{=}\sum_{k=1}^{m}\zeta_k{\circ}\bar\zeta_k$, where $a{\circ}b{=}\frac{1}{2}(a{*}b{+}b{*}a)$. First of all, note that (29) and the comment given in § 3.3.1 give the following:

Proposition 3.9 *In generic ordered expressions* $e_*^{t\rho_*^2}{=}\prod_{k=1}^{m}{*}e_*^{t\zeta_k\circ\bar\zeta_k}$ *and* $\int_0^\infty e_*^{-t\rho_*^2}dt$ *is defined to give an inverse of* ρ_*^2*. Moreover, the Laplace transform gives*

$$\rho_*^{-1}{=}\sqrt[*]{\rho_*^2}^{-1}{=}\frac{1}{\sqrt{\pi}}\int_0^\infty\frac{1}{\sqrt{t}}e_*^{-t\rho_*^2}dt$$

As ρ_*^2 has the property $E_s(\rho_*^2){=}e^{2s}\rho_*^2$, changing variables in the integration gives easily $E_s\rho_*^{-1}{=}e^{-s}\rho_*^{-1}$.

E_s-invariant elements are generated by

$$\nu{*}\rho_*^{-2}, \quad \rho_*^{-1}{*}\zeta_k, \quad \bar\zeta_k{*}\rho_*^{-1}, \quad k = 1{\sim}m.$$

We denote the generated algebra by $\mathcal{A}(S^{2m-1})$ in the space $\mathcal{E}_{2+}(\mathbb{C}^{2m})$.

For simplicity we denote

$$\mu{=}\nu{*}\rho_*^{-2}, \quad \xi_k{=}\rho_*^{-1}{*}\zeta_k, \quad \bar\xi_k{=}\bar\zeta_k{*}\rho_*^{-1}, \quad k = 1{\sim}m.$$

Although they are not hermitian, we see easily

$$[\mu^{-1},\xi_k]{=}\xi_k, \quad [\mu^{-1},\bar\xi_k]{=}{-}\bar\xi_k, \quad \sum_{k=1}^{m}\xi_k{*}\bar\xi_k{=}1{-}2m\mu. \tag{35}$$

It follows that $e^{it\mathrm{ad}(\mu^{-1})}(\xi_k){=}e^{it}\xi_k$, $e^{it\mathrm{ad}(\mu^{-1})}(\bar\xi_k){=}e^{-it}\bar\xi_k$, and $\mathrm{Ad}(e_*^{it\mu^{-1}})$ causes an S^1-action on $\mathcal{A}(S^{2m-1})$. Hence S^1-invariant elements form a subalgebra $\mathcal{A}(P_{m-1}(\mathbb{C}))$ generated by

$$\mu, \quad \xi_k{*}\bar\xi_\ell, \quad \bar\xi_k{*}\xi_\ell, \quad 1\leq k,\ell\leq m.$$

Furthermore, as $\rho_*^2{*}\zeta_k{=}\zeta_k{*}(\rho_*^2{+}2\nu)$, $\rho_*^2{*}\bar\zeta_k{=}\zeta_k{*}(\rho_*^2{-}2\nu)$, we have

$$\rho_*^{-1}{*}\zeta_k{=}\frac{1}{\sqrt{\pi}}\int_0^\infty\frac{1}{\sqrt{t}}e_*^{-t\rho_*^2}{*}\zeta_k dt{=}\zeta_k{*}\frac{1}{\sqrt{\pi}}\int_0^\infty\frac{1}{\sqrt{t}}e_*^{-t(\rho_*^2+2\nu)}dt{=}\zeta_k{*}\frac{1}{\sqrt{\rho_*^2{+}2\nu}}.$$

Using $\sqrt{\rho_*^2{+}2\nu}{=}\rho_*{*}\sqrt{1{+}2\mu}{=}\sqrt{1{+}2\mu}{*}\rho_*$, we have

$$\rho_*^{-1}{*}\zeta_k{=}\zeta_k{*}\rho_*^{-1}{*}\sqrt{1{+}2\mu}^{-1}$$

Similar calculation gives

$$\bar\zeta_k{*}\rho_*^{-1}{=}\sqrt{1{+}2\mu}^{-1}{*}\rho_*^{-1}{*}\bar\zeta_k, \quad \rho_*^{-1}{*}\bar\zeta_k{=}\bar\zeta_k{*}\rho_*^{-1}{*}\sqrt{1{-}2\mu}^{-1}, \quad \zeta_k{*}\rho_*^{-1}{=}\sqrt{1{-}2\mu}^{-1}{*}\rho_*^{-1}{*}\zeta_k.$$

By using first two identities (without using $\sqrt{1{-}2\mu}$) we have that

$$[\xi_k, \xi_\ell]{=}0{=}[\bar{\xi}_k, \bar{\xi}_\ell], \quad [\xi_k, \bar{\xi}_\ell]{=}-\frac{1}{1{+}2\mu}{*}(2\mu\delta_{k\ell}{+}\xi_k{*}\bar{\xi}_\ell) \tag{36}$$

where we used the fact that

$$[\mu^{-1}, \zeta_k{*}\bar{\zeta}_\ell]{=}\nu^{-1}{*}[\rho_*^2, \zeta_k{*}\bar{\zeta}_\ell]{=}0, \quad [\sqrt{1{+}2\mu}, \rho]{=}0{=}[\sqrt{1{+}2\mu}, \zeta_k{*}\bar{\zeta}_\ell].$$

Proposition 3.10 *The algebra $\mathcal{A}(S^{2m-1})$ generated by $\{\mu,\ \xi_k,\ \bar{\xi}_k; k = 1{\sim}m\}$ is a contact Weyl algebra such that $\mathrm{ad}(i\mu^{-1})$ gives an S^1-action on $\mathcal{A}(S^{2m-1})$. Commutation relations are given by* (35) *and* (36)*.*

For every non-negative integer $n{\in}\mathbb{N}$, we see $[\mu^{-1}, \xi^n_{i*}]{=}n\xi^n_{i*}$, $[\mu^{-1}, \bar{\xi}^n_{i*}]{=}-n\bar{\xi}^n_{i*}$. Hence $\mathcal{A}(S^{2m-1})$ has the eigenspace decomposition of $\mathrm{ad}(\mu^{-1})$

$$\mathcal{A}(S^{2m-1}){=}\sum_{\ell\in\mathbb{Z}}\oplus\mathcal{A}_\ell(S^{2m-1}), \quad \mathcal{A}_\ell(S^{2m-1}){*}\mathcal{A}_{\ell'}(S^{2m-1}){=}\mathcal{A}_{\ell+\ell'}(S^{2m-1}), \quad \ell, \ell' \in \mathbb{Z}.$$

The subalgebra $\mathcal{A}_0(S^{2m-1}) = \mathcal{A}(P_{m-1}(\mathbb{C}))$ may be viewed as the algebra obtained by the reduction procedure.

3.4.1 Complex vacuum representation

Computations in this subsection execute in generic ordered expressions and the suffix $:\ :_{_K}$ will be omitted. Here we use the vacuum (32) Hence we have $\bar{\xi}_i{*}\widetilde{\varpi}_{\mathbb{C}}{=}0$, $\sum_{k=1}^m \xi_k{*}\bar{\xi}_k{*}\widetilde{\varpi}_{\mathbb{C}}{=}(1{-}2m\mu){*}\widetilde{\varpi}_{\mathbb{C}}{=}0$.

As $\rho_*^2{*}\widetilde{\varpi}_{\mathbb{C}}{=}2m\nu\widetilde{\varpi}_{\mathbb{C}}$, we have

$$\rho_*^{-1}{*}\widetilde{\varpi}_{\mathbb{C}}{=}\frac{1}{\sqrt{\pi}}\int_0^\infty \frac{1}{\sqrt{t}}e^{-t2m\nu}dt\widetilde{\varpi}_{\mathbb{C}}{=}\frac{1}{\sqrt{2m\nu}}\widetilde{\varpi}_{\mathbb{C}}, \quad \sqrt{\mu}{*}\widetilde{\varpi}_{\mathbb{C}}{=}\frac{1}{\sqrt{2m}}\widetilde{\varpi}_{\mathbb{C}}.$$

$$\rho_*^2{*}h(\boldsymbol{\zeta}){*}\widetilde{\varpi}_{\mathbb{C}}{=}\sum_k(\nu\zeta_k\partial_k{+}2\nu)h(\boldsymbol{\zeta}){*}\widetilde{\varpi}_{\mathbb{C}}, \quad \mu^{-1}{*}h(\boldsymbol{\zeta}){*}\widetilde{\varpi}_{\mathbb{C}}{=}\sum_k(\zeta_k\partial_k{+}2)h(\boldsymbol{\zeta}){*}\widetilde{\varpi}_{\mathbb{C}}.$$

In particular, for every polynomial $p_k(\boldsymbol{\zeta})$ of degree k

$$\mu^{-1}{*}p_k(\boldsymbol{\zeta}){*}\widetilde{\varpi}_{\mathbb{C}}{=}(k{+}2m)p_k(\boldsymbol{\zeta}){*}\widetilde{\varpi}_{\mathbb{C}}, \quad \mu{*}p_k(\boldsymbol{\zeta}){*}\widetilde{\varpi}_{\mathbb{C}}{=}\frac{1}{k{+}2m}p_k(\boldsymbol{\zeta}){*}\widetilde{\varpi}_{\mathbb{C}}.$$

As $e_*^{s\sum\zeta_i\circ\bar{\zeta}_i}{*}p_k(\boldsymbol{\zeta}){*}\widetilde{\varpi}_{\mathbb{C}}{=}e^{s(k+2m\nu)}p_k(\boldsymbol{\zeta}){*}\widetilde{\varpi}_{\mathbb{C}}$, we have

$$\xi_i{*}p_k(\boldsymbol{\zeta}){*}\widetilde{\varpi}_{\mathbb{C}}{=}\frac{1}{\sqrt{k{+}1{+}2m\nu}}\zeta_i p_k(\boldsymbol{\zeta}){*}\widetilde{\varpi}_{\mathbb{C}}, \quad \bar{\xi}_i{*}p_k(\boldsymbol{\zeta}){*}\widetilde{\varpi}_{\mathbb{C}}{=}\frac{1}{\sqrt{k{+}2m\nu}}\partial_i p_k(\boldsymbol{\zeta}){*}\widetilde{\varpi}_{\mathbb{C}}.$$

Hence, there is no canonical conjugate of μ^{-1}.

One can use $\mathbb{C}[\boldsymbol{\xi}]{*}\widetilde{\varpi}_{\mathbb{C}}$ as the representation space. Then

$$\mu^{-1}{*}p_k(\boldsymbol{\xi}){*}\widetilde{\varpi}_{\mathbb{C}}{=}(k{+}2m)p_k(\boldsymbol{\xi}){*}\widetilde{\varpi}_{\mathbb{C}},$$

The formula for $\bar{\xi}_i{*}p_k(\boldsymbol{\xi}){*}\widetilde{\varpi}_{\mathbb{C}}$ is complicated.

3.4.2 Localizations joining divisors

Consider now localizations by joining divisors, say

$$(\bar{\zeta}_k{*}\zeta_k)_+^{-1}, \quad k=1{\sim}m,$$

where $(\bar{\zeta}_k{*}\zeta_k)_+^{-1}{=}\int_0^\infty e_*^{-t\bar{\zeta}_k*\zeta_k}dt$. $\bar{\zeta}^{\bullet}_k{=}\zeta_k{*}(\bar{\zeta}_k{*}\zeta_k)_+^{-1}$ is a left-inverse of ζ i.e. $\bar{\zeta}{*}\bar{\zeta}^{\bullet}_k{=}1$, but

$$\bar{\zeta}^{\bullet}_k{*}\bar{\zeta}{=}1{-}\lim_{t\to-\infty} e_*^{-t\zeta_k*\bar{\zeta}_k} \quad \text{(partial complex vacuum)}.$$

Recalling the generator system (23), we consider a new generator system for E_t-invariant elements in generic ordered expressions

$$\nu{*}(\zeta_1{*}(\bar{\zeta}_1{*}\zeta_1)_+^{-1})^2,\ \frac{1}{2}\zeta_1^2{*}(\bar{\zeta}_1{*}\zeta_1)_+^{-1},\ \ \zeta_k{*}\zeta_1{*}(\bar{\zeta}_1{*}\zeta_1)_+^{-1}, \quad \zeta_1{*}(\bar{\zeta}_1{*}\zeta_1)_+^{-1}{*}\bar{\zeta}_k, \quad k=2{\sim}m$$

For simplicity, denote these by

$$\mu,\ \tau,\ \eta_k,\ \bar{\eta}_k, \quad k=2{\sim}m,$$

and let $\mathcal{A}$ be the algebra generated by these in $\mathcal{E}_{2+}(\mathbb{C}^{2m})$.

By using

$$[\zeta_1,\zeta_1{*}(\bar{\zeta}_1{*}\zeta_1)_+^{-1}]{=}\zeta_1{*}[\zeta_1,(\bar{\zeta}_1{*}\zeta_1)_+^{-1}]{=}{-}\zeta_1{*}(\bar{\zeta}_1{*}\zeta_1)_+^{-1}{*}[\zeta_1,\bar{\zeta}_1{*}\zeta_1]{*}(\bar{\zeta}_1{*}\zeta_1)_+^{-1}{=}2\nu{*}(\zeta_1{*}(\bar{\zeta}_1{*}\zeta_1)_+^{-1})^2,$$

we easily see that

$$[\mu,\tau]{=}2\mu^2, \quad [\tau,\eta_k]{=}\mu{*}\eta_k, \quad [\tau,\bar{\eta}_k]{=}\mu{*}\bar{\eta}_k, \quad [\eta_k,\bar{\eta}_\ell]{=}\mu\delta_{k\ell}, \quad \text{others are } 0.$$

Hence, this system generates a contact Weyl algebra with a canonical conjugate of μ^{-1}. Moreover, one may join ζ_1 to $\mathcal{A}$ and $\mathcal{A}[\zeta_1]$ is a μ-regulated algebra. However, just as in Proposition 3.6, there is no standard vacuum ϖ such that $\zeta_1{*}\varpi{=}0$ or $\varpi{*}\zeta_1{=}0$ holds. This is because $\varpi{*}\zeta_1{=}0$ gives $\varpi{*}\mathcal{A}{=}\{0\}$ and $\zeta_1{*}\varpi{=}0$ gives $\mu{*}\varpi{=}\infty$.

4 The case embedded Heisenberg algebra

We saw that the semi-boundedness of μ^{-1} suffers the existence of its canonical conjugate, and the localizations by joining a divisor or an $*$-inverse of some element defined by integrals suffers the existence of standard abstract vacuum.

Recall now the embedding, given in § 3.2.2 Theorem 3.3 into $\mathcal{E}_{1+}(\mathbb{C}^{2m+2})$. This may be viewed as a local normal form of the system mentioned in the previous subsection. In this case, the element $\rho^2{=}e_*^{2y_0}$ is brought from the outside of original Heisenberg algebra $\mathcal{H}_{2m}[\mu]$. Namely, $\mu^{-1}{=}\hbar^{-1}e_*^{2y_0}$ is an *external energy variable*. As $\mu{=}\hbar e_*^{-2y_0}$, and $\tau{=}\frac{1}{2}(e_*^{-2y_0}{*}x_0{+}x_0{*}e_*^{-2y_0})$ is a canonical conjugate of μ^{-1}, τ may be viewed as the external time.

In the extended Weyl algebra generated by x_0, y_0, we denote the vacuum

$$\varpi_{00}{=}\lim_{t\to-\infty} e^{\frac{1}{2}t}e_*^{t\frac{1}{i\hbar}x_0\circ y_0}.$$

Note first that ϖ_{00} in not in $\mathcal{E}_{1+}(\mathbb{C}^{2m+2})$, but in $\mathcal{E}_{2+}(\mathbb{C}^{2m+2})$.

The abstract vacuum $\widetilde{\varpi}_s(\mathcal{H}){=}e_*^{-is\tau}{*}\widetilde{\varpi}(L_0)$ given in (34) will be called the **Heisenberg vacuum** after showing the remarkable property that $e_*^{-is\tau}{*}\widetilde{\varpi}(L_0)=0$ for $\frac{1}{1+2\hbar s}\leq 0$ in the later section (cf. Theorem 4.2). Using $[\mu^{-1}, e_*^{-is\tau}]{=}s\hbar e_*^{-is\tau}$ and $\mu^{-1}{*}\varpi_{00}{=}\frac{1}{\hbar}\varpi_{00}$, we have

$$\mu{*}e_*^{-is\tau}{*}\widetilde{\varpi}(L_0){=}\begin{cases}\frac{\hbar}{1+2\hbar s}e_*^{-is\tau}{*}\widetilde{\varpi}(L_0), & \frac{1}{1+2\hbar s}>0\\ \qquad 0, & \frac{1}{1+2\hbar s}\leq 0\end{cases}$$

We take the representation space as

$$C^{\infty}(\mathbb{R}_{s>s_0}{\times}\mathbb{R}^m){*}\widetilde{\varpi}_s(\mathcal{H}),\quad s_0=-\frac{1}{2\hbar}.$$

Then, we see $\tilde{u}_k$ is represented as a multiplication $\tilde{u}_k{*}f(s,\tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_s(\mathcal{H}){=}(\tilde{u}_k{*}f(s,\tilde{\boldsymbol{u}})){*}\widetilde{\varpi}_s(\mathcal{H})$, and

$$\mu{*}f(s,\tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_s(\mathcal{H}){=}\frac{\hbar}{1{+}2\hbar s}f(s,\tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_s(\mathcal{H}),\quad \tilde{v}_k{*}f(s,\tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_s(\mathcal{H}){=}-\frac{\hbar i}{1{+}2\hbar s}\partial_k f(s,\tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_s(\mathcal{H}).$$

μ^{-1} and τ are canonical conjugate pair. τ is represented by a similar form to $X=\frac{1}{i\mu}\sum_{k=1}^{m}\tilde{u}_k{*}\tilde{v}_k$

$$\begin{aligned}X{*}f(s,\tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_s(\mathcal{H}) &= \frac{\hbar i}{1{+}2s\hbar}\sum_{k=1}^{m}\tilde{u}_k\partial_{\tilde{u}_k}f(s,\tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_s(\mathcal{H})\\ i\tau{*}f(s,\tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_s(\mathcal{H}) &= \frac{\hbar}{1{+}2s\hbar}\sum_{k=1}^{m}\tilde{u}_k\partial_{\tilde{u}_k}f(s,\tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_s(\mathcal{H}){+}f(s,\tilde{\boldsymbol{u}}){*}i\partial_s\widetilde{\varpi}_s(\mathcal{H})\end{aligned}\tag{37}$$

In the representation space μ behaves as $\frac{i}{1+2\hbar s}$. Tending $s\to\infty$ gives $\mu\to 0$. This gives a classical limit. The most remarkable fact is that $\mathrm{Ad}(e_*^{it\tau})$ **is well-defined only for** $t\geq 0$ (cf. Proposition 4.4). Hence,

$$e_*^{it\tau}{*}f(s){*}\widetilde{\varpi}_s(\mathcal{H}){=}f(s){*}\widetilde{\varpi}_{s+t}(\mathcal{H})$$

for only $t{>}0$. In general,

$$e_*^{it\mu^{-1}\tau}{*}f(s,\boldsymbol{u}){*}\widetilde{\varpi}_s(\mathcal{H}){=}f(s,E_{a(s)t}(\boldsymbol{u}))\widetilde{\varpi}_{s+t}(\mathcal{H}),\quad t\geq 0,\quad a(s){=}\frac{\hbar}{1{+}2s\hbar}$$

Vacuum representation with respect to $\widetilde{\varpi}_{y_0}(\overline{L}_0)$

On the other hand, as $(x_0{+}i\hbar){*}e_*^{y_0}{*}\overline{\varpi}_{00}{=}e_*^{y_0}{*}x_0{*}\overline{\varpi}_{00}$, the abstract vacuum (33) satisfies $\tau{*}\widetilde{\varpi}_{y_0}(\overline{L}_0){=}0$ in generic ordered expression, and hence

$$e_*^{it\tau}{*}\widetilde{\varpi}_{y_0}(\overline{L}_0){=}\widetilde{\varpi}_{y_0}(\overline{L}_0),\quad t{\in}\mathbb{C}.$$

As $[i\tau,\mu^{-1}]{=}[i\tau,\mu^{-1}]{=}2$, we have

$$[i\tau,f(\mu)]{=}-2\mu^2f'(\mu),\quad [i\tau,\sqrt{\mu}]{=}\frac{1}{\sqrt{\mu}}$$

It follows

$$i\tau{*}f(\mu^{-1}){*}\widetilde{\varpi}_{y_0}(\overline{L}_0){=}[i\hat{\tau}, f(\mu^{-1})]{*}\widetilde{\varpi}_{y_0}(\overline{L}_0){=}2f'(\mu^{-1}){*}\widetilde{\varpi}_{y_0}(\overline{L}_0).$$

On the other hand, as $\tilde{u}_k{=}\frac{\sqrt{\hbar}}{\sqrt{\mu}}{*}x_k$, we have $[i\hat{\tau}, \tilde{u}_k]{=}\mu{*}\tilde{u}_k$, hence

$$i\hat{\tau}{*}f(\tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_{y_0}(\overline{L}_0){=}\mu{*}\sum_k \tilde{u}_k\partial_{\tilde{u}_k}f(\tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_{y_0}(\overline{L}_0).$$

Let $Hol(\mathbb{C}^m)$ be the space of holomorphic functions of $\tilde{u}_1\cdots, \tilde{u}_m$. We set the representation space by

$$Hol(\mathbb{C}^{m+1})[\mu, \mu^{-1}]]{*}\widetilde{\varpi}_{y_0}(\overline{L}_0).$$

Then, we see $\tilde{u}_k$ and μ^{-1} are represented as multiplications and

$$\tau{*}f(\mu^{-1}, \tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_{y_0}(\overline{L}_0){=}i\Big(2\partial_{\mu^{-1}}{+}\mu{*}\sum_k \tilde{u}_k\partial_{\tilde{u}_k}\Big)f(\mu^{-1}, \tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_{y_0}(\overline{L}_0). \tag{38}$$

In particular, considering the equation with initial data $\mu^{-1}{*}\widetilde{\varpi}_{y_0}(\overline{L}_0)$,

$$\frac{d}{dt}e_*^{it\tau}{*}\mu^{-1}{*}\widetilde{\varpi}_{y_0}(\overline{L}_0){=}e_*^{it\tau}{*}[\hat{\tau}, \mu^{-1}]{*}\widetilde{\varpi}_{y_0}(\overline{L}_0){=}2{*}\widetilde{\varpi}_{y_0}(\overline{L}_0),$$

we obtain

$$e_*^{it\tau}{*}\mu^{-1}{*}\widetilde{\varpi}_{y_0}(\overline{L}_0){=}(\mu^{-1}{+}2t){*}\widetilde{\varpi}_{y_0}(\overline{L}_0), \quad t{\in}\mathbb{C}.$$

In general

$$e_*^{it\mu^{-1}\tau}{*}f(\mu^{-1}, \tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_{y_0}(\overline{L}_0){=}f(\mu^{-1}(1{+}2t), E_t(\tilde{\boldsymbol{u}})){*}\widetilde{\varpi}_{y_0}(\overline{L}_0)$$

This gives also

$$e_*^{it\tau}{*}\mu{*}\widetilde{\varpi}_{y_0}(\overline{L}_0){=}\frac{1}{\mu^{-1}{+}2t}{*}\widetilde{\varpi}_{y_0}(\overline{L}_0){=}\frac{\mu}{1{+}2t\mu}{*}\widetilde{\varpi}_{y_0}(\overline{L}_0), \quad t{\in}\mathbb{C}.$$

One may take $C^\infty(\mathbb{R}^m)[\mu^{-1}, \mu]]{*}\widetilde{\varpi}_{y_0}(\overline{L}_0)$ as the representation space, where μ and $\tilde{u}_k$ are represented by multiplication operator, and

$$e_*^{it\mu^{-1}{*}\tau}{*}f(\mu, \tilde{\boldsymbol{u}}){*}\widetilde{\varpi}_{y_0}(\overline{L}_0){=}f\big(\frac{1}{1{+}2t\mu}, E_t(\tilde{\boldsymbol{u}})\big){*}\widetilde{\varpi}_{y_0}(\overline{L}_0).$$

Here, to avoid singularities, we have to treat μ in a formal μ-regulated algebra by setting

$$\frac{1}{1{+}2t\mu}{=}(1{-}2t\mu{+}(2t\mu)^2{+}(2t\mu)^3{-}\cdots.$$

4.1 The star-exponential functions $e_*^{it\tau}{*}F$, $F{*}e_*^{-it\tau}$

To give the proof of Theorem 4.2, we start with a little general setting. Now, recalling

$$:\tau:_0{=}e^{-2y_0}x_0, \quad x_0{*}e^{-2y_0}{=}e^{-2y_0}{*}(x_0{+}2i\hbar)$$

we consider the $*$-exponential function $:e_*^{ite^{-2y_0}*(x_0+i\hbar)}:_{_K}$, or $*$-the product $:e_*^{ite^{-2y_0}*(x_0+i\hbar)}{*}F:_{_K}$.

The equation for these are

$$\begin{aligned} &\frac{d}{dt}F_t = :(ie_*^{-2y_0}{*}(x_0{+}i\hbar)):_{_K}{*_{_K}}F_t, \quad F_0 = :F:_{_K} \\ &\frac{d}{dt}\hat{F}_t = \hat{F}_t{*_{_K}}:(ie_*^{-2y_0}{*}(x_0{+}i\hbar)):_{_K}, \quad \hat{F}_0 = :F:_{_K}. \end{aligned} \tag{39}$$

As the Hermitian structure is defined, applying the anti-automorphism $a \to \bar{a}$ to the first one to get

$$\frac{d}{dt}\overline{F}_t = -\overline{F}_t{*_{_K}}:(ie_*^{-2y_0}{*}(x_0{+}i\hbar)):_{_K}, \quad \overline{F}_0 = :\overline{F}:_{_K}.$$

Minding the uniqueness of the real analytic solution, we may assume that $F_t = F_t(x_0, y_0)$ and all computations carry out within these two variables.

In what follows, all computations execute under the Weyl (K=0) ordered expression. If the expression parameter K is restricted to $K{=}0$ (Weyl ordered expression), we denote $:\ :_0$ in such computation. Since

$$(x_0{+}i\hbar){*_0}F_t(x_0,y_0){=}(x_0{+}i\hbar)F_t(x_0,y_0){-}\frac{i\hbar}{2}\partial_{y_0}F_t(x_0,y_0), \quad (x_0{+}i\hbar){*_{_{\hat{K}}}}F_t(x_0,y_0){=}(x_0{+}i\hbar)F_t(x_0,y_0),$$

and

$$e^{-2y_0} *_0 f(x_0,y_0) = e^{-2y_0}f(x_0{-}i\hbar, y_0), \quad e^{-2y_0} *_{_{\hat{K}}} f(x_0,y_0) = e^{-2y_0}f(x_0{-}2i\hbar, y_0),$$

we see that (39) turns out to be

$$\frac{d}{dt}F_t(x_0,y_0) = ie^{-2y_0}x_0F_t(x_0{-}i\hbar, y_0) + \frac{\hbar}{2}e^{-2y_0}\partial_{y_0}F_t(x_0{-}i\hbar, y_0), \tag{40}$$

with initial condition F_0.

Putting $F_t(x_0,y_0) = G_t(x_0,y_0)e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}$, the equation (40) becomes

$$\frac{d}{dt}G_t(x_0,y_0) = \frac{\hbar}{2}e^{-4y_0}\partial_{y_0}G_t(x_0{-}i\hbar, y_0), \quad G_0 = F_0(x_0,y_0)e^{-\frac{2}{i\hbar}(x_0+i\hbar)y_0}. \tag{41}$$

(41) is not a differential equation, but a differential-difference equation.

The case $G_t(x_0, y_0)$ is periodic.

If G_t is restricted to periodic functions $G_t(x_0{-}i\hbar, y_0) = G_t(x_0, y_0)$, then (41) turns out to be

$$\big(\partial_t - \frac{\hbar}{2}e^{-4y_0}\partial_{y_0}\big)G_t(x_0,y_0) = 0,$$

and the solution with the initial data is given by

$$G_t(x_0,y_0) = \varphi\big(x_0, 2\hbar t{+}e^{4y_0}\big), \quad G_0(x_0,y_0) = \varphi\big(x_0, e^{4y_0}\big)$$

by using every holomorphic function $\varphi(u,v)$ such that $\varphi(u{+}i\hbar,v)=\varphi(u,v)$. Thus the initial function $G_0(x_0,y_0)$ must satisfy the periodical conditions $G_0(x_0{-}i\hbar,y_0){=}G_0(x_0,y_0)$. This is equivalent with $F(x_0{-}i\hbar,y_0){=}F(x_0,y_0)e^{-2y_0}$.

Hence $:e^{it\tau}:_0$ is not defined by itself but it is defined only in the form $:e^{it\tau}{*}F_*:_0$. That is in the form of solutions of equations written under the Weyl ordered expression with the initial data F_*.

For any $\varphi(x_0,y_0)$ satisfying $\varphi(x_0{-}i\hbar,y_0){=}\varphi(x_0,y_0)$ (e.g. $\varphi(y_0)$), we set $F{=}\varphi(x_0,e^{4y_0})e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}$. Then, we see

$$:e_*^{it\tau}:_0{*_0}\varphi(x_0,e^{4y_0})e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}{=}\varphi(x_0,2\hbar t{+}e^{4y_0})e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}. \tag{42}$$

Thus, if $t{\in}\mathbb{C}$, then $\varphi(x_0,y_0)$ must be entire w.r.t. y_0. Recalling

$$:\overline{\varpi}_{00}:_0{=}2e^{\frac{2}{i\hbar}x_0y_0}, \quad :f(y_0){*}\overline{\varpi}_{00}:_0{=}f(2y_0)2e^{\frac{2}{i\hbar}x_0y_0},$$

we see $e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}{=}:e^{y_0}{*}\overline{\varpi}_{00}:_0$.

In particular, restricting φ to a entire function $\varphi(z)$ and noting that $\varphi(e^{2y_0}){=}:\varphi(e_*^{2y_0}):_0$, (42) is written as

$$:e_*^{it\tau}{*}\varphi(e_*^{2y_0}){*}e_*^{y_0}{*}\overline{\varpi}_{00}:_0{=}:\varphi(e_*^{2y_0}{+}2\hbar t){*}e_*^{y_0}{*}\overline{\varpi}_{00}:_0.$$

Although the computation was done under the Weyl ordered expression, one may write this without suffix $:\ :_0$, that is,

$$e_*^{it\tau}{*}\varphi(e_*^{2y_0}){*}e_*^{y_0}{*}\overline{\varpi}_{00}=\varphi(e_*^{2y_0}{+}2\hbar t){*}e_*^{y_0}{*}\overline{\varpi}_{00}, \tag{43}$$

as all terms in r.h.s. can be intertwined to generic expressions.

Recall that $e_*^{y_0}{*}\overline{\varpi}_{00}$ is an abstract vacuum such that

$$\overline{\varpi}_{00}{*}y_0{=}0, \quad (x_0{+}i\hbar){*}e_*^{y_0}{*}\overline{\varpi}_{00}=0, \quad i\tau{*}e_*^{y_0}{*}\overline{\varpi}_{00}=0.$$

Hence, (43) may be written as

$$e_*^{it\tau}{*}\varphi(e_*^{2y_0}){*}e_*^{-it\tau}{*}e_*^{y_0}{*}\overline{\varpi}_{00}=\mathrm{Ad}(e_*^{it\tau})\varphi(e_*^{2y_0}){*}e_*^{y_0}{*}\overline{\varpi}_{00}{=}\varphi(e_*^{2y_0}{+}2\hbar t){*}e_*^{y_0}{*}\overline{\varpi}_{00}, \quad t{\in}\mathbb{C}.$$

On the other hand, it is easy to see that $[i\tau,e_*^{2y_0}]{=}2\hbar$, $[i\tau,[i\tau,e_*^{2y_0}]]{=}0,\cdots$. This gives

$$\exp t\mathrm{ad}(i\tau)\varphi(e_*^{2y_0}){=}\varphi(e_*^{2y_0}{+}2\hbar t), \quad t{\in}\mathbb{C}$$

for every entire function φ.

Applying these to $e_*^{-2y_0}$, we have

$$e_*^{it\tau}{*}e_*^{-2y_0}{*}e_*^{y_0}{*}\overline{\varpi}_{00}=(e_*^{2y_0}{+}2\hbar t)^{-1}{*}e_*^{y_0}{*}\overline{\varpi}_{00}{=}e_*^{-2y_0}\frac{1}{1{+}2\hbar te_*^{-2y_0}}{*}e_*^{y_0}{*}\overline{\varpi}_{00}.$$

$$\exp t\mathrm{ad}(i\tau)(e_*^{-2y_0}){=}e_*^{-2y_0}\frac{1}{1{+}2\hbar te_*^{-2y_0}}, \quad t{\in}\mathbb{C}.$$

This is the same to the previous case $\widetilde{\varpi}_{y_0}(\overline{L}_0)$.

Fourier transform expressions

Next we want to remove the periodical condition $G_0(x_0{-}i\hbar, y_0) = G_0(x_0, y_0)$. Instead, we assume that t is a real variable and $G_t(x_0, y_0)$ is written as

$$G_t(x_0, y_0) = \int_{-\infty}^{\infty} a(t, \xi, y_0) e^{i\xi\, x_0} {\mathchar'26\mkern-12mu d}\xi, \quad G_0(x_0, y_0) = \int_{-\infty}^{\infty} a(0, \xi, y_0) e^{i\xi\, x_0} {\mathchar'26\mkern-12mu d}\xi$$

by using tempered distribution $a(t, \xi, y_0)$ of compact support with respect to ξ.

If x_0, y_0 in (44) are treated as real variables, then the compactness of the support is removed, but $G_t(x_0{-}i\hbar, y_0)$ must be viewed as the shift of ξ

$$G_t(x_0{-}i\hbar, y_0) = \int_{-\infty}^{\infty} a(t, \xi, y_0) e^{i\xi(x_0-i\hbar)} {\mathchar'26\mkern-12mu d}\xi = \int_{-\infty}^{\infty} a(t, \xi, y_0) e^{\hbar\xi} e^{i\xi x_0} {\mathchar'26\mkern-12mu d}\xi.$$

Precisely, $a(t, \xi, y_0)$ is regarded for every (t, y_0) as an $\mathcal{S}'$-valued function $a(t, y_0)(\xi)$ for every test function $\psi(\xi) \in \mathcal{S}$. It is remarkable in the Weyl ordered expression that the equation (41) is changed into the differential equation

$$\big(e^{-\hbar\xi}\partial_t - \frac{\hbar}{2} e^{-4y_0}\partial_{y_0}\big) a(t, \xi, y_0) = 0 \tag{44}$$

involving ξ as a parameter, i.e.

$$\int \Big(e^{-\hbar\xi}\partial_t a(t, y_0)(\xi) - \frac{\hbar}{2} e^{-4y_0}\partial_{y_0} a(t, y_0)(\xi)\Big)\psi(\xi) {\mathchar'26\mkern-12mu d}\xi = 0.$$

(44) shows that $a(t, \xi, y_0)$ is constant along the real vector field $e^{-\hbar\xi}\partial_t - \frac{\hbar}{2} e^{-4y_0}\partial_{y_0}$ on the (y_0, t)-surface. That is constant along the integral curves of this vector field. Hence, the solution is

$$a(t, \xi, y_0) = \phi(\xi, 2\hbar t{+}e^{-\hbar\xi} e^{4y_0}), \quad a(0, \xi, y_0) = \phi(\xi, e^{-\hbar\xi} e^{4y_0}) \tag{45}$$

by using a function $\phi(\xi, \eta)$ on $\mathbb{R}^2$ such that $f(\xi) = \phi(\xi, 2\hbar t{+}e^{-\hbar\xi} e^{4y_0})$ is a tempered distribution of compact support. If we allow x_0 to be a real variable, then the condition of compact support is relaxed simply as follows:

Condition 1. $h(\eta) = \phi(\xi, \eta)$ is a $\mathcal{S}'$-valued C^∞ function such that for every η, and for every (t, y_0),

$$\phi(\xi, 2\hbar t{+}e^{-\hbar\xi} e^{4y_0})$$

is also a tempered distribution w.r.t. ξ.

Lemma 4.1 *If $\phi(\xi, \eta)$ is a C^∞ function such that $|\partial_\eta^n \phi(\xi, \eta)\eta^n| < \infty$ for every n, then Condition 1 is fulfilled.*

Proof As

$$\frac{d}{d\xi}\phi(\xi, 2\hbar t{+}e^{-\hbar\xi} e^{4y_0}) {=} (\partial_\xi \phi(\xi, 2\hbar t{+}e^{-\hbar\xi} e^{4y_0}) {+} \hbar e^{-\hbar\xi} e^{4y_0} \partial_\eta \phi(\xi, 2\hbar t{+}e^{-\hbar\xi} e^{4y_0})).$$

Repeating this it is enough to show the boundedness of

$$(\hbar e^{-\hbar\xi} e^{4y_0})^n \partial_\eta^n \phi(\xi, 2\hbar t{+}e^{-\hbar\xi} e^{4y_0}))$$

This is easily checked. □

So far, the variable y_0 has been regarded as a complex variable, but Condition 1 allows x_0 to be a real variable. Such an unbalanced situation yields the following:

Theorem 4.1 *If the initial condition* $a(0,\xi,y_0)$ *has not periodic property* $a(0,\xi,y_0){=}a(0,\xi,y_0{\pm}\frac{1}{2}\pi i)$*, then there is no solution of* (39)*. This periodical condition is equivalent with* $G(x_0,y_0{\pm}\frac{1}{2}\pi i){=}G(x_0,y_0)$*, and with* $F(x_0,y_0{\pm}\frac{1}{2}\pi i){=}F(x_0,y_0)e^{\frac{\pi}{\hbar}x_0}$*.*

In particular, the complex $*$*-exponential function* $e_*^{z\tau}$ *cannot exist by itself. It exists only in the form* $e_*^{z\tau}{*}F$ *with initial function* F *with a quasi-periodical property* $F(x_0,y_0{+}\pi i){=}F(x_0,y_0)e^{\frac{\pi}{\hbar}x_0}$*.*

As t and ξ are real variables, it is natural to allow y_0 to be a real variable. Then, (44) can be solved without any restriction.

To simplify the argument, we put several assumptions in what follows:

Assumption. $\hbar$ is a positive constant. t, ξ and y_0 are real variables. Under such restrictions, $:e_*^{it\tau}:_0$ may be expressed as a function $F_t(x_0,y_0)$ of real variables.

In real variable expressions, it occurs a strange phenomenon. Fix ξ arbitrarily. If we set $\eta{=}2\hbar t{+}e^{-\hbar\xi}e^{4y_0}$, then $a(t,\xi,y_0)$ is constant on the curve $\eta{=}2\hbar t{+}e^{-\hbar\xi}e^{4y_0}$. Thus, if $\eta\leq 0$, then $\eta{-}e^{-\hbar\xi}e^{4y_0}{=}2\hbar t$ is negative. Such a curve does not cross the initial line $t=0$. As the initial data is given at $t=0$, the initial data must be given only when $\eta{>}0$ by the form

$$a(0,\xi,y_0)=\phi(\xi,\eta),\quad \eta=e^{-\hbar\xi}e^{4y_0}\quad i.e.\ 4y_0=\hbar\xi{+}\log\eta. \tag{46}$$


Graphs of $-2\hbar t=e^{-\hbar\xi+4y_0}{-}\eta$
$\eta=0$
$\eta<0$
$\eta>0$
The data on dotted line disappear on the line $t=0$
The deta on the line $t=0$ is used only on the right hand side
The line $t=0$
y_0

Another word, we have

Proposition 4.1 *If* $a(0,\xi,y_0){=}a(0,\xi,\frac{1}{4}(\hbar\xi{+}\log\eta))=0$ *for* $\eta>0$*, then* $a(t,\xi,y_0)=0$ *for* $\forall t\geq 0$*, that is,* $G_t(x_0,y_0)=0$ *for* $\forall t\geq 0$*.*

Given $\phi(\xi,\eta)$, the data at $t=0$ is $\phi(\xi,e^{-\hbar\xi+4y_0})$ given by using only the values on $\eta>0$, and the data at $t=t_0$ is $\phi(\xi,e^{-\hbar\xi+4y_0}{+}2\hbar t_0)$. Some data at $t=0$ disappears in the time $t_0{>}0$, and some data on $\eta{<}0$ is used at $t_0{<}0$.
Another word **one may choose arbitrary function on the domain** $\eta<0$. For (39) **the uniqueness does not hold in the direction** $t_0{<}0$.

In spite of this, we note the next

Proposition 4.2 *If one sets* $\phi(\xi,\eta)=0$ *for* $\eta\leq 0$*, namely one does not use the data for* $\eta\leq 0$*, then the solution of* (39) *is uniquely determined by the data at* $t=0$*.*

Now, start with an initial ($t{=}0$) data $a(0,\xi,y_0){=}a(0,\xi,\frac{1}{4}(\hbar\xi{+}\log\eta))$, and take the unique solution $a(s,\xi,\frac{1}{4}(\hbar\xi{+}\log\eta))$ at s ($s>0$). This is uniquely determined but the data at $0{<}\eta{<}2\hbar s$ is not used in the data at $t=s$. Now, want to restart using the data at $t{=}s$ as the initial data and go back to the line $t=0$. Then the original data at $t{=}0$ is not recovered. This procedure violates the exponential law, that is, the associativity of the one parameter product formula.

Now, integrating the result in Proposition 4.1 by ξ, we have

Proposition 4.3 *For an arbitrary tempered distribution* $\phi(\xi,\eta)$ *w.r.t* ξ *satisfying Condition 1 and* $\phi(\xi,\eta)=0$ *for* $\eta>0$*, the integral* $G_t(x_0,y_0){=}\int_{\mathbb{R}}\phi(\xi,2\hbar t{+}e^{-\hbar\xi+4y_0})e^{i\xi x_0}{\mathchar'26\mkern-12mu d}\xi$ *satisfies* $G_t(x_0,y_0){=}0$ *for* $t{\geq}0$*. Therefore,* $F_t(x_0,y_0){=}G_t(x_0,y_0)e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}$ *satisfies the equation* (39)*, but* $F_t(x_0,y_0){=}0$ *for* $t{\geq}0$*.*

If one wants to use the value at $t=t_0>0$ as the initial data, then $-2\hbar(t+t_0)=e^{-\hbar\xi+4y_0}+\eta$ is used instead of $-2\hbar t=e^{-\hbar\xi+4y_0}+\eta$, where nonvanishing data remains.

Since the uniqueness ensured only by setting $\phi(\xi,\eta)=0$ on the area $\eta<0$, we see

Proposition 4.4 *The adjoint operator* $\mathrm{Ad}(e_*^{it\tau})F=e_*^{it\tau}*F*e_*^{-it\tau}$ *is defined only for $t\geq 0$ by using*

$$e_*^{it\tau}*\overline{(e_*^{it\tau}*\overline{F})},\quad t\geq 0.$$

Let $\mathcal{I}(0)$ be the space of all compactly supported tempered distribution $\phi(\xi,\eta)$ w.r.t. ξ. Every $\phi(\xi,\eta)\in\mathcal{I}(0)$ satisfies Condition 1. We set $\hat{g}_a(\xi,y_0)=\phi(\xi,2\hbar t+e^{-\hbar\xi+4y_0})$ for $t\in\mathbb{R}$ and

$$:g_t(x_0,y_0):_0=\int_{\mathbb{R}}\hat{g}_t(\xi,y_0)e^{i\xi x_0}đ\xi.$$

Then, the integral

$$F_t(x_0,y_0)=\int_{\mathbb{R}}\hat{g}_t(\xi,\frac{1}{4}(\hbar\xi+\log(e^{-\hbar\xi+4y_0}+2\hbar t))e^{i\xi x_0}đ\xi\, e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}$$

is the solution of (40) uniquely determined for the "future" direction by the data given at $t=a$. Hence one may write

$$F_t(x_0,y_0)=:e_*^{it\tau}:_0*_0F_0(x_0,y_0),\quad t\geq 0.$$

By the uniqueness gives the following:

Proposition 4.5 *The exponential law* $:e_*^{i(s+t)\tau}:_0*_0F_0(x_0,y_0)=:e_*^{is\tau}*e_*^{it\tau}:_0*_0F_0(x_0,y_0)$ *holds for $s,t>0$.*

The next one may sound surprising:

Proposition 4.6 *If the initial $\phi(\xi,\eta)\in\mathcal{I}(0)$ is compactly supported w.r.t. (ξ,η), then there is $c>0$ such that $:e_*^{it\tau}:_0*_0F_0(x_0,y_0)=0$ for $t\geq c$. $F_0(x_0,y_0)$ may be viewed as* **an element of a finite time span of life**.

Note As $i\tau$ is skew-hermitian, we have the primitive conservation law:

$$\frac{d}{dt}\overline{F}_t*F_t=0. \tag{47}$$

The above result does not against the conservation law (47). This implies simply $\overline{F}_t*F_t=0$. Note also $\overline{\varpi}_{00}*\varpi_{00}=0$ (cf. [19]).

Recall that (42) shows $:e_*^{it\tau}:_0*_0e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}=e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}$.

4.1.1 Solutions for the initial value $e^{a\frac{1}{i\hbar}x_0y_0}$

By virtue of the restriction to the real variables, one can obtain the solution for many initial values.

Consider next $:e_*^{it\tau}:_0*_0e^{a\frac{1}{i\hbar}x_0y_0}$ for general $a\in\mathbb{R}$. Noting that $e^{(a-2)\frac{1}{i\hbar}x_0y_0}=\int_{\mathbb{R}}\delta(\xi+\frac{a-2}{\hbar}y_0)e^{i\xi x_0}đ\xi$, we have the initial data

$$e^{a\frac{1}{i\hbar}x_0y_0}=\int e^{-2y_0}\delta(\xi+\frac{a-2}{\hbar}y_0)e^{i\xi x_0}đ\xi e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}.$$

Hence the initial data $\phi(\xi,\eta)$ in (46) is given by

$$\phi(\xi,\eta)=e^{-2y_0}\delta(\xi{+}\frac{a{-}2}{\hbar}y_0){=}e^{-\frac{1}{2}(\hbar\xi+\log\eta)}\delta((1{+}\frac{a{-}2}{4})\xi{+}\frac{a{-}2}{4\hbar}\log\eta),$$

plugging $y_0=\frac{1}{4}(\hbar\xi{+}\log(\eta))$. As $\delta(x)$ is supported only on $x=0$, the support of $\phi(\xi,\eta)$ is given as the curve $(\frac{a+2}{4})\xi{+}\frac{a-2}{4\hbar}\log\eta=0$ in the (ξ,η) plane:


η
The support of
$\phi(\xi,\eta)$ for $\frac{a+2}{a-2}<0$
1
ξ
any function below this line
η
The support of
$\phi(\xi,\eta)$ for $a=2$
0
ξ
any function below this line

But this gives $\phi(\xi,\eta)$ only on “half” area $\eta>0$.

To obtain the uniquness, we set $\phi(\xi,\eta)=0$ for $\eta{<}0$. General solution is obtained by adding arbitrary $\phi(\xi,\eta)$ supported on $\eta<0$.

By this convention, the solution is obtained by replacing η by $2\hbar t{+}e^{-\hbar\xi+4y_0}$ providing $2\hbar t{+}e^{-\hbar\xi+4y_0}{>}0$ and $\eta{=}0$ if $2\hbar t{+}e^{-\hbar\xi+4y_0}{\leq}0$.

$$:e_*^{it\tau}:_0*_0e^{a\frac{1}{i\hbar}x_0y_0}=\int e^{-\frac{1}{2}(\hbar\xi+log(e^{4y_0-\hbar\xi}+2\hbar t))}\delta\Big((\frac{a{+}2}{4})\xi{+}\frac{a{-}2}{4\hbar}\log(e^{4y_0-\hbar\xi}{+}2\hbar t)\Big)e^{i\xi x_0}\bar{d}\xi\ e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}. \tag{48}$$

The case $a{=}-2$. This is the case that we want for the proof of Theorem 4.2, where $\phi(\xi,\eta)=e^{-\frac{1}{2}(\hbar\xi+\log\eta)}\delta(-\frac{1}{\hbar}\log(\eta))$.



That is, the support of $\phi(\xi,\eta)$ is the line $\eta=1$ in the (ξ,η) plane. Setting $\xi'=-\frac{1}{\hbar}\log(e^{-\hbar\xi+4y_0}{+}2\hbar t)$, we see that if $1{-}2\hbar t\leq 0$, then ξ' cannot be 0. Thus, $\delta(\xi')=0$ for $1{-}2\hbar t\leq 0$. The solution may be written as

$$:e_*^{it\tau}:_0*_0e^{-2\frac{1}{i\hbar}x_0y_0}=\begin{cases} 0 & 2\hbar t\geq 1\\ e^{-\frac{\hbar}{2}(\xi-\xi')}\int\delta(\xi')e^{i\xi x_0}\bar{d}\xi\ e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0} & 2\hbar t<1\end{cases}$$

$\xi'=0$ is at $\hbar\xi=4y_0{-}\log(1-2\hbar t)$, i.e. $e^{-\frac{1}{2}\hbar\xi}|_{\xi'=0}=\sqrt{1{-}2\hbar t}\,e^{-2y_0}$. It follows

$$\frac{d\xi'}{d\xi}\Big|_{\xi'=0}=\frac{e^{-\hbar\xi}e^{4y_0}}{e^{-\hbar\xi}e^{4y_0}{+}2\hbar t}=1{-}2\hbar t.$$

Thus,

$$:e_*^{it\tau}:_0*_0e^{-2\frac{1}{i\hbar}x_0y_0}{=}\begin{cases} 0 & 2\hbar t\geq 1\\ \frac{1}{\sqrt{1-2\hbar t}}e^{-i\frac{x_0}{\hbar}\log(1-2\hbar t)}e^{-\frac{2}{i\hbar}x_0y_0} & 2\hbar t{<}1\end{cases} \tag{49}$$

Recall the vacuum $:\varpi_{00}:_0=2e^{-2\frac{1}{i\hbar}x_0y_0}$ and the formula $:f(x_0):_0{*_0}:\varpi_{00}:_0 = f(2x_0):\varpi_{00}:_0$. As the r.h.s. is defined on generic ordered expression, Proposition 3.3 gives that (49) is rewritten without $:\ :_0$ as a rather surprising form

Theorem 4.2 *In generic ordered expression, we have*

$$e_*^{it\tau}*\frac{1}{2}\varpi_{00}=\begin{cases} 0 & 2\hbar t \geq 1 \\ \frac{1}{\sqrt{1-2\hbar t}}e_*^{\frac{x_0}{2i\hbar}\log(1-2\hbar t)}*\frac{1}{2}\varpi_{00} & 2\hbar t<1. \end{cases}$$

It is not hard to see that

$$\sum_{n=0}^{\infty}\frac{1}{n!}(\frac{t}{\hbar})^n\tau_*^n*\varpi_{00}=\sum_{n=0}^{\infty}\frac{1}{n!}(\frac{t}{\hbar})^n\frac{i\hbar}{x_0}(\frac{x_0}{i\hbar})_{n+1}*\varpi_{00}$$

without μ.

Hence (49) is the "time evolution" of the vacuum. This is real analytic for $t<\frac{1}{2\hbar}$ and the exponential law holds on this domain. However, Proposition 4.3 shows that solutions are not unique on $\eta = e^{4y_0}+2\hbar t<0$ depending on arbitrary functions.

4.1.2 The case that initial date is $\overline{\varpi}_{00}$

Consider now the case a=2. Note first that $:\overline{\varpi}_{00}:_0 = 2e^{\frac{2}{i\hbar}x_0y_0}$. Hence this case gives the vacuum representation w.r.t. $\overline{\varpi}_{00}$. If this is the case, the property of the delta function gives

$$:e_*^{it\tau}*\frac{1}{2}\overline{\varpi}_{00}:_0 = e^{-\frac{1}{2}log(e^{4y_0}+2\hbar t)}e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}=\frac{1}{\sqrt{e^{4y_0}+2\hbar t}}e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}=\frac{1}{\sqrt{1+2\hbar te^{-4y_0}}}e^{\frac{2}{i\hbar}x_0y_0} \tag{50}$$

This is the case $\varphi(z)=(\sqrt{z})^{-1}$ in (**??**). It looks that there is a double branched singularities. Here we have to consider the treatment of $\sqrt{\ }$.

By noting that $:(\sqrt[*]{1+2\hbar te_*^{-2y_0}})_*^{-1}:_0=\frac{1}{\sqrt{1+2\hbar te^{-2y_0}}}$ and $f(y_0){*_0}:\overline{\varpi}_{00}:_0=f(2y_0):\overline{\varpi}_{00}$, (50) is rewritten as

$$:e_*^{it\tau}*:\overline{\varpi}_{00}:_0 = :\Big(\sqrt[*]{1+2\hbar te_*^{-2y_0}}\Big)_*^{-1}:_0{*_0}:\overline{\varpi}_{00}:_0,$$

Compare this with the result (**??**). As real variables are considered, we have to set $\phi(\xi,\eta) = 0$ for $\eta = e^{4y_0}+2\hbar t < 0$ by the comment between Proposition 4.2 and Proposition 4.3 to make the solution unique. Thus, the solution might be written by the discontinuous function

$$:e_*^{it\tau}*\frac{1}{2}\overline{\varpi}_{00}:_0 = \Big(\frac{1}{\sqrt{1+2\hbar te^{-4y_0}}}\Big)_{re}e^{\frac{2}{i\hbar}x_0y_0}, \quad \Big(\frac{1}{\sqrt{1+2\hbar te^{-4y_0}}}\Big)_{re} = \begin{cases} \frac{1}{\sqrt{1+2\hbar te^{-4y_0}}} & e^{4y_0}+2\hbar t > 0 \\ 0 & e^{4y_0}+2\hbar t < 0. \end{cases}$$

Note. What is moving by the equation (39) is not a point, but a function. Hence the singular set $2\hbar t+e^{4y_0}=0$ is *not* a singular point of the equation (39). It is only that the initial smooth function is changed into a function involving a singular point depending on t.

However, if we think these under formal power series of $\hbar$, then $\frac{1}{\sqrt{1+2\hbar te^{-4y_0}}}$ makes sense as a formal power series of $\hbar e^{-4y_0}$. Indeed, using $x_0{*}e_*^{-2ny_0}{=}e_*^{-2ny_0}{*}(x_0{+}2ni\hbar)$, we have in generic ordered expression that

$$\sum_{k=0}^{\infty}\frac{(it)^k}{k!}{:}\tau_*^k{*}\overline{\varpi}_{00}{:}_K{=}{:}\Big(\sqrt[*]{1{+}2\hbar te_*^{-2y_0}}\Big)_*^{-1}{*}\overline{\varpi}_{00}{:}_K{=}{:}\frac{1}{\sqrt[*]{1{+}2t\mu}}{*}\overline{\varpi}_{00}{:}_K \tag{51}$$

where the r.h.s. stands for the Taylor series of μ.

Proposition 4.7 *Although $e_*^{it\tau}{*}e_*^{y_0}{*}\overline{\varpi}_{00}{=}e_*^{y_0}{*}\overline{\varpi}_{00}$ holds in generic ordered expression, $e_*^{it\tau}{*}\overline{\varpi}_{00}$ is defined only in a formal μ-regulated algebra.*

On the other hand, taking the hermitian conjugate by using the Hermitian structure, we have

$$\varpi_{00}{*}e_*^{y_0}{*}e_*^{-it\tau}=\varpi_{00}{*}e_*^{y_0}. \tag{52}$$

The combination of (52) and (49) gives under generic ordered expression that

$$\begin{aligned}&(\varpi_{00}{*}e_*^{y_0}{*}e_*^{-it\tau}){*}\varpi_{00}{=}(\varpi_{00}{*}e_*^{y_0}){*}\varpi_{00}\\ &\varpi_{00}{*}e_*^{y_0}{*}(e_*^{-it\tau}{*}\varpi_{00}){=}\varpi_{00}{*}e_*^{y_0}{*}\big(\frac{1}{\sqrt{1{+}2\hbar t}}{*}\varpi_{00}\big),\end{aligned} \tag{53}$$

hence the associativity is broken. This is not a contradiction, because these are elements of $\mathcal{E}_{2+}(\mathbb{C}^{2m})$.
The reason seems that $\varpi_{00}{*}e_*^{y_0}$ is a vacuum different from ϖ_{00}.

On the contrary, in the formal level we see the associativity holds: The combination of the hermitian conjugate of (51) and (49) gives by formal power series of μ

$$\begin{aligned}{:}(\varpi_{00}{*}e_*^{-it\tau}){*}\varpi_{00}{:}_0 &= \Big(2e^{-\frac{2}{i\hbar}x_0y_0}\frac{1}{\sqrt{1{+}2\hbar te^{-4y_0}}}\Big){*}_0{:}\varpi_{00}{:}_0{=}\Big({:}\varpi_{00}{*}(\sqrt[*]{1{+}2\hbar te^{-2y_0}})_*^{-1}{:}_0\Big){*}_0{:}\varpi_{00}{:}_0\\ {:}\varpi_{00}{*}(e_*^{-it\tau}{*}\varpi_{00}){:} &= {:}\varpi_{00}{:}_0{*}_0\Big(\frac{1}{\sqrt{1{+}2\hbar t}}e^{-i\frac{x_0}{\hbar}\log(1+2\hbar t)}2e^{-\frac{2}{i\hbar}x_0y_0}\Big)\\ &{=}{:}\varpi_{00}{:}_0{*}_0\Big(\frac{1}{\sqrt{1{+}2\hbar t}}e^{-i\frac{x_0}{2\hbar}\log(1+2\hbar t)}{*}_0{:}\varpi_{00}{:}_0\Big)\end{aligned}$$

by using ${:}\varpi_{00}{*}y_0{:}_0{=}{:}\varpi_{00}{:}_0 2y_0$, ${:}x_0{*}\varpi_{00}{:}_0{=}{:}\varpi_{00}{:}_0 2x_0$. At a first glance, these violate the associativity, but in fact as $y_0{*}\varpi_{00}{=}0{=}\varpi_{00}{*}x_0$, we have the associativity

$${:}(\varpi_{00}{*}e_*^{-it\tau}){*}\varpi_{00}{:}_0{=}\frac{1}{\sqrt{1{+}2\hbar t}}{:}\varpi_{00}{:}_0{=}{:}\varpi_{00}{*}(e_*^{-it\tau}{*}\varpi_{00}){:}_0, \quad 2\hbar t{\geq}-1. \tag{54}$$

4.1.3 Solution for the initial value 1

It is interesting to investigate the solution with the initial value 1, but in this subsection, by technical reason, all computation are restricted under the Weyl ordered ($K=0$) expression.

Note that the initial condition $F_0 = 1$ corresponds to $\phi(\xi, y_0) = e^{-2y_0}\delta(\xi - \frac{2}{\hbar}y_0)$, as

$$G_0(x_0, y_0) = e^{-\frac{2}{i\hbar}(x_0+i\hbar)y_0} = \int_{-\infty}^{\infty} e^{-2y_0}\delta(\xi - \frac{2}{\hbar}y_0)e^{i\xi\, x_0}\bar{d}\xi.$$

Since this is the case $a = 0$, (48) gives that the solution obtained by putting $\phi(\xi, \eta) = 0$ for $\eta < 0$ is

$$:e_*^{it\tau}:_0 = \int_{-\infty}^{\infty} e^{-\frac{1}{2}(\hbar\xi + \log(e^{-\hbar\xi+4y_0}+2\hbar t))}\delta\Big(\frac{1}{2}\xi - \frac{1}{2\hbar}\log(e^{-\hbar\xi+4y_0} + 2\hbar t)\Big)e^{i\xi\, x_0}\bar{d}\xi\, e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}.$$

This is the unique real analytic solution and $e^{-\hbar\xi+4y_0}+2\hbar t > 0$ is satisfied.

Note that $\delta\Big(\frac{1}{2}\xi - \frac{1}{2\hbar}\log(e^{-\hbar\xi+4y_0}+2\hbar t)\Big)$ is supported only on the point $\frac{1}{2}\xi - \frac{1}{2\hbar}\log(e^{-\hbar\xi+4y_0}+2\hbar t)=0$. Moreover,

$$\xi' = \frac{1}{2}\xi - \frac{1}{2\hbar}\log(e^{-\hbar\xi+4y_0}+2\hbar t)$$

is easily inverted as

$$\xi = \frac{1}{\hbar}\log\Big(\sqrt{(\hbar t e^{4\hbar\xi'})^2+e^{4y_0}e^{2\hbar\xi'}} + \hbar t e^{2\hbar\xi'}\Big).$$

Hence

$$e^{\hbar\xi}\Big|_{\xi'=0} = e^{4y_0-\hbar\xi}\Big|_{\xi'=0}+2\hbar t = \sqrt{(\hbar t)^2+e^{4y_0}}+\hbar t, \qquad \frac{d\xi'}{d\xi} = \frac{\sqrt{(\hbar t)^2+e^{4y_0}}}{\sqrt{(\hbar t)^2+e^{4y_0}}+\hbar t}.$$

Thus, the change of variable of integration gives

$$:e_*^{it\tau}:_0 = \int_{-\infty}^{\infty} e^{-\frac{1}{2}(\hbar\xi + \log(e^{4y_0-\hbar\xi}+2\hbar t))}\delta(\xi')e^{i\xi x_0}\frac{d\xi}{d\xi'}\bar{d}\xi'\, e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0} = \int_{-\infty}^{\infty}\Big(e^{-\hbar\xi}e^{i\xi x_0}\frac{d\xi}{d\xi'}\Big)\Big|_{\xi'=0}\bar{d}\xi'\, e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}$$

as $\delta(\xi')$ is supported only on $\xi' = 0$. Hence we obtain

$$:e_*^{it\tau}:_0 = :e_*^{ite_*^{-2y_0}*(x_0+ih)}:_0 = \frac{1}{\sqrt{(\hbar t)^2+e^{4y_0}}}e^{\frac{ix_0}{\hbar}\log(\sqrt{(\hbar t)^2+e^{4y_0}}+\hbar t)}e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0}$$

As $\sqrt{(\hbar t)^2+e^{4y_0}}+\hbar t > 0$ always, this is the unique real analytic solution with initial condition 1. We denote this by

$$\begin{aligned} :\exp_*(it\tau):_0 &= \frac{1}{\sqrt{(\hbar t)^2+e^{4y_0}}}e^{\frac{ix_0}{\hbar}\log(\sqrt{(\hbar t)^2+e^{4y_0}}+\hbar t)}e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0} \\ &= \frac{1}{\sqrt{1+(e^{-2y_0}\hbar t)^2}}e^{\frac{ix_0}{\hbar}\log(\sqrt{1+(e^{-2y_0}\hbar t)^2}+e^{-2y_0}\hbar t)}, \quad t \in \mathbb{R} \end{aligned} \tag{55}$$

This may be rewritten as

$$:\exp_*(it\tau):_0 = \frac{1}{\sqrt{1+(\mu t)^2}}e^{\frac{ix_0}{\hbar}\log(\sqrt{1+(\mu t)^2}+\mu t)}.$$

Note also that

$$(\sqrt{1+(\mu t)^2}+\mu t)(\sqrt{1+(\mu t)^2}-\mu t)=1, \quad (\sqrt{1+(\mu t)^2}+\mu t)+(\sqrt{1+(\mu t)^2}-\mu t)=\sqrt{1+(\mu t)^2}.$$

This expression has no singularity and it is easy to check

$$\overline{:\exp_*(it\tau):_0} = :\exp_*(-it\tau):_0.$$

Note that $:i\tau:_0 = ie^{-2y_0}x_0$ to check $:\frac{d}{dt}\Big|_{t=0}e_*^{it\tau}:_0 = :i\tau:_0$.

If an arbitrary $\tilde{\phi}(\eta)$ supported on the domain $\eta < 0$ is considered, then

$$:e_*^{it\tau}:_0 = :\exp_*(it\tau):_0 + \int_{-\infty}^{\infty} \tilde{\phi}(2\hbar t{+}e^{-\hbar\xi}e^{4y_0})e^{i\xi x_0} đ\xi \, e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0} \tag{56}$$

and solutions are not unique in the past direction.

By these calculations and the uniqueness for future direction, we must have the exponential law

$$:\exp_*(is\tau):_0{*_0}:\exp_*(it\tau):_0 = :\exp_*(i(s{+}t)\tau):_0, \quad s,t \geq 0$$

holds, and if $*_0F$ in the r.h.s. is welldefined then

$$:\exp_*(it\tau):_0{*_0}F = \frac{1}{\sqrt{(\hbar t)^2{+}e^{4y_0}}} e^{\frac{ix_0}{\hbar}\log(\sqrt{(\hbar t)^2+e^{4y_0}}+\hbar t)} e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0} {*_0}F$$

is the solution of initial data F.

Computational conjecture.

By the observation above, one must have the identities

$$\begin{aligned} :e_*^{-it\tau}{*}\frac{1}{2}\varpi_{00}:_0 = :e_*^{-it\tau}:_0{*_0}e^{-\frac{2}{i\hbar}x_0y_0} &= \frac{1}{\sqrt{(\hbar t)^2{+}e^{4y_0}}} e^{\frac{ix_0}{\hbar}\log(\sqrt{(\hbar t)^2+e^{4y_0}}-\hbar t)} e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0} {*_0}e^{-\frac{2}{i\hbar}x_0y_0} \\ &= \frac{1}{\sqrt{1{+}2\hbar t}} e^{-i\frac{x_0}{\hbar}\log(1+2\hbar t)} e^{-\frac{2}{i\hbar}x_0y_0}, \quad -1{<}2\hbar t. \end{aligned}$$

But it is not easy to check these identities by direct calculations.

It seems possible to prove directly that in some restricted expressions

$$\begin{aligned} :e_*^{it\tau}{*}\frac{1}{2}\overline{\varpi}_{00}:_K &= \frac{1}{\sqrt{(\hbar t)^2{+}e^{4y_0}}} e^{\frac{ix_0}{\hbar}\log(\sqrt{(\hbar t)^2+e^{4y_0}}+\hbar t)} e^{\frac{2}{i\hbar}(x_0+i\hbar)y_0} {*_K} e^{\frac{2}{i\hbar}x_0y_0} \\ &= \frac{1}{\sqrt{1{+}2\hbar t e^{-4y_0}}} e^{\frac{2}{i\hbar}x_0y_0}, \quad e^{4y_0}{+}2\hbar t > 0. \end{aligned}$$

5 Berezin algebra embedded in the extended Weyl algebra

As it is well known, Berezin operators give operator representations of algebras which may be viewed as the quantization of the structure of the Poincar'e unit disk, where the central element corresponding to $\hbar$ is represented by an operator. Note that the notion of $*$-product is not used, but as it is naturally expected, the $*$-product that we used in this note is redefined by using the operator product. In this section, we give an embedding of Berezin algebra into the extended Weyl algebra by tracing [4].

5.1 Algebra on the unit disk D.

To define Berezin operators we first remark the following:

Lemma 5.1 *For every (anti-) holomorphic function g on D with suitable growth condition so that the integrals converge, we have for $s>-1$*

$$\frac{(s+1)i}{2\pi}\iint_D g(v)(1-v\bar{v})^s dv d\bar{v} = g(0) = \frac{(s+1)i}{2\pi}\iint_D g(\bar{v})(1-v\bar{v})^s dv d\bar{v},$$

Proof By the Taylor expansion of $g(v)$ at 0 and noting the integral vanishes by the integration $\int e^{ik\theta}d\theta$ for $k\neq 0$, we have only to compute as follows:

$$\frac{i}{2}\iint_D (1-v\bar{v})^s dv d\bar{v} = 2\pi\int_0^1 (1-r^2)^s r dr = \pi\int_0^1 (1-t)^s dt = \frac{\pi}{s+1}.$$

□

By the definition of the beta function, we see also the following:

$$\frac{i}{2}\iint_D (v\bar{v})^m(1-v\bar{v})^s dv d\bar{v}=2\pi\int_0^1 r^{2m}(1-r^2)^s r dr=\pi B(m+1,s+1), \quad \mathrm{Re}\, m+1,\ \mathrm{Re}\, s+1>0 \tag{57}$$

Thus, setting

$$\Phi_s(w,v,\bar{v}) = \frac{1}{\pi}\sum_{m=0}^{\infty}\frac{1}{B(m+1,s+1)}(1-v\bar{v})^s(w\bar{v})^m = \frac{s+1}{\pi}\frac{(1-v\bar{v})^s}{(1-w\bar{v})^{s+2}}$$

we have the identity

$$f(w) = \int_D f(v)\Phi_s(w,v,\bar{v})dv d\bar{v} = \frac{s+1}{\pi}\int_D f(v)\left(\frac{1-v\bar{v}}{1-w\bar{v}}\right)^{s+2} dm(v,\bar{v}) \tag{58}$$

where $dm(v,\bar{v}) = \frac{i}{2(1-v\bar{v})^2}dv d\bar{v}$ is the volume element on the Poincaré unit disk.

Let $\mathcal{H}^s(D)$ be the space of all holomorphic functions on D such that

$$\int_D |f|^2(1-v\bar{v})^{s+2}dm(v,\bar{v}) < \infty.$$

It is known that $e^m_{(s)} = (\frac{\Gamma(s+m+2)}{\pi\Gamma(m+1)\Gamma(s+1)})^{1/2}w^m$ form an orthonormal basis of $\mathcal{H}^s(D)$. Note that $s>-1$ is crucial to obtain Hilbert spaces, and no vacuum appears here.

Consider the linear space bundle and the space of all smooth sections

$$\coprod_{s>-1}\mathcal{H}^s(D), \quad \mathit{\Gamma}(\coprod_{s>-1}\mathcal{H}^s(D)).$$

For a function $a(w,\bar{w})$ on D, the Berezin operator $P_s(a)$ on $\mathcal{H}^s(D)$ is defined by

$$(P_s(a)f)(w) = \frac{(s+1)i}{2\pi}\int_D a(v,\bar{v})f(v)\Phi_s(w,v,\bar{v})dv d\bar{v} = \frac{(s+1)}{\pi}\int_D a(v,\bar{v})f(v)\left(\frac{1-v\bar{v}}{1-w\bar{v}}\right)^{s+2} dm(v,\bar{v}). \tag{59}$$

Let $P_s(a)e^m_{(s)} = \sum_k A_{m,k}e^k_{(s)}$. Then, we have

$$A_{m,k} = \frac{1}{\pi}\int_D a(v,\bar{v})v^m\bar{v}^k(1-v\bar{v})^s dv d\bar{v}(\tfrac{\Gamma(s+m+2)}{\Gamma(m+1)\Gamma(s+1)})^{1/2}(\tfrac{\Gamma(s+k+2)}{\Gamma(k+1)\Gamma(s+1)})^{1/2} \tag{60}$$

Hence the operator norm $||P_s(a)||$ is given by

$$||P_s(a)||^2 = (s+1)^2 \iint a(v,\bar{v})\overline{a(w,\bar{w})}\left(\frac{1-v\bar{v}}{1-v\bar{w}}\frac{1-w\bar{w}}{1-w\bar{v}}\right)^{s+2} dm(v,\bar{v})dm(w,\bar{w})$$

It is easy to see that

$$\begin{aligned}(P_s(\phi(w))f)(w) &= \int_D \phi(v)f(v)\Phi_s(w,v,\bar{v})dvd\bar{v} = \phi(w)f(w),\\ (P_s(\frac{\bar{w}}{1-w\bar{w}})f)(w) &= \int_D \frac{\bar{v}}{1-v\bar{v}}f(v)\Phi_s(w,v,\bar{v})dvd\bar{v} = \frac{1}{s}f'(w)\end{aligned} \tag{61}$$

The second one is defined only for $s>0$. We define the product $* = *_s$ by the operator product, i.e.

$$P_s(a*b) = P_s(a)P_s(b).$$

For a holomorphic function f on the unit disk D, let $f(w) = \sum_k a_k w^k$. We have then setting $\pi B(s+1,m+1) = \pi\frac{\Gamma(m+1)\Gamma(s+1)}{\Gamma(s+m+2)}$,

$$\begin{aligned}(P_s(\bar{w}^l))f(w) &= \sum_{k,m}\int_D a_k\bar{v}^l v^k\frac{1}{\pi B(s+1,m+1)}(w\bar{v})^m(1-\bar{v}v)^s dvd\bar{v}\\ &= \sum_m \int_D a_{m+l}w^m\frac{1}{\pi B(s+1,m+1)}(v\bar{v})^{m+l}(1-\bar{v}v)^s dvd\bar{v}\\ &= \sum_m a_{m+l}w^m\frac{B(s+1,m+l+1)}{B(s+1,m+1)} = \sum_m a_{m+l}w^m\frac{(m+l)\cdots(m+1)}{(s+m+l+1)\cdots(s+m+2)}\end{aligned} \tag{62}$$

This is a bounded operator such that $(P_s(\bar{w}^l))w^m = 0$ for $0 \le m \le l-1$. Using (62), we have

$$\begin{aligned}(P_s(\bar{w}*w)f)(w) &= (P_s(\bar{w})P_s(w))f(w)\\ &= \sum_{k,m}\int_D a_k\bar{v}v^{k+1}\frac{1}{\pi B(s+1,m+1)}(w\bar{v})^m(1-\bar{v}v)^s dvd\bar{v}\\ &= \sum_{k\ge 0}\int_D a_k w^k(\bar{v}v)^{k+1}(1-\bar{v}v)^s\frac{1}{\pi B(s+1,k+1)} = \sum_k a_k\frac{B(s+1,k+2)}{B(s+1,k+1)}w^k\\ &= \sum_k a_k\frac{k+1}{s+k+2}w^k\end{aligned} \tag{63}$$

Hence, $P_s(\bar{w}*w)$ is a bounded diagonal operator, which is invertible within bounded operators. Note also that

$$P_s(\bar{w}*w) = P_s(\bar{w}\cdot w) \neq P_s(w*\bar{w}), \quad P_s((\bar{w}\cdot w)^k) \neq P_s((\bar{w}\cdot w)^k_*).$$

Denote by $(\bar{w}*w)_*^{-1}$ the inverse of $\bar{w}*w$. Then, $w^\bullet=(\bar{w}*w)_*^{-1}*\bar{w}$ (resp. $\bar{w}^\circ=w*(\bar{w}*w)_*^{-1}$) is a left (resp. right) inverse of w (resp. $\bar{w}$):

$$\begin{aligned}&((\bar{w}{*}w)_*^{-1}*\bar{w})*w=1, \quad w*((\bar{w}{*}w)_*^{-1}*\bar{w})=1-\varpi_0,\\ &\bar{w}*(w*(\bar{w}{*}w)_*^{-1})=1, \quad (w*(\bar{w}{*}w)_*^{-1})*\bar{w}=1-\varpi_0\end{aligned} \tag{64}$$

where ϖ_0 is the element corresponding to the projection operator given by $P_s(\varpi_0)(f)=f(0)$.

By (63) we have that

$$(P_s(1-\bar{w}{*}w))^{-1}f(w)=(P_s((1-\bar{w}{*}w)_*^{-1})f)(w)=\sum_k a_k w^k\frac{s+k+2}{s+1}$$

This is an unbounded operator. Using this, we have

$$P_s((1-\bar{w}{*}w)_*^{-1}*\bar{w})f(w)=\frac{1}{s+1}f'(w), \quad s\geq 0. \tag{65}$$

Note that $P_s((1-\bar{w}{*}w)_*^{-1}*\bar{w})=\frac{s}{s+1}P_s(\frac{\bar{w}}{1-w\bar{w}})$.

By (64), we see that $w*(\bar{w}{*}w)^{-1}*(1-\bar{w}{*}w)=w*(\bar{w}{*}w)^{-1}-w$ is a right inverse of $(1-\bar{w}{*}w)_*^{-1}*\bar{w}$. Indeed, we see that

$$P_s(w*(\bar{w}{*}w)^{-1}-w)=(s+1)\int_0^w dw.$$

For each $s>0$, let $\mathbb{D}_{(s)}$ be the algebra of operators generated by $w,\bar{w}$, $(\bar{w}*w)^{-1}$, $(1-\bar{w}*w)^{-1}$, $(1-w*\bar{w})^{-1}$, $\hbar$.

Proposition 5.1 *$w^*=(1-\bar{w}{*}w)_*^{-1}*\bar{w}$ is a canonical conjugate of w i.e. $[w^*,w]{=}\frac{1}{s+1}$, and*

$$[w,\bar{w}]=-\frac{1}{s+1}(1-w{*}\bar{w})*(1-\bar{w}{*}w).$$

Hence the Berezin algebra $\mathbb{D}_{(s)}$ may be viewed as a deformation quantization of the Kähler structure defined on the unit disk D by the Poisson bracket $\{w,\bar{w}\}{=}-\frac{1}{s+1}(1-w\bar{w})^2$.

Proof Note that $-\frac{1}{s+1}=[w,(1-\bar{w}{*}w)^{-1}*\bar{w}]$. Hence we have

$$-\frac{1}{s+1}=(1-\bar{w}{*}w)_*^{-1}*[w,\bar{w}]*w*(1-\bar{w}{*}w)_*^{-1}*\bar{w}+(1-\bar{w}{*}w)^{-1}*[w,\bar{w}]$$

By the bumping lemma, the right hand side is

$$(1-\bar{w}{*}w)_*^{-1}*[w,\bar{w}]*((1-w{*}\bar{w})_*^{-1}*w{*}\bar{w}+1)=(1-\bar{w}{*}w)_*^{-1}*[w,\bar{w}]*(1-w{*}\bar{w})_*^{-1}$$

This give the result. □

Note also that ϖ_0 can be obtained by using (61) as

$$\varpi_0=\lim_{t\to\infty}e_*^{-tw*\frac{\bar{w}}{1-\bar{w}w}}=\lim_{t\to\infty}e^{-\frac{t}{s}w\partial_w}.$$

Note here that the algebra above is given directly as operators. Indeed, this is a deformation quantization of Poincaré disk in the notion that mentioned in § 2. However, in the previous sections, the concrete calculation is done by using $*_K$-product by using an expression parameter K.

5.2 Embeddings into the extended Weyl algebra

On the other hand, one can easily make a deformation quantization of the Kähler structure on the Poincaré disk D in $\mathcal{E}_{2+}(\mathbb{C}^3$. Let ν, x, y be a generator of $\mathcal{H}_2[\nu]$. We set $z{=}x{+}iy$, $\bar{z}{=}x{-}iy$. Using the $*$-exponential function $e_*^{t\bar{z}*z}$ we see that

$$\sqrt[*]{\bar{z}*z}^{\,-1}, \quad \sqrt[*]{1{+}\bar{z}*z}^{\,-1}$$

are welldefined in $\mathcal{E}_{2+}(\mathbb{C}^3)$. We set as follows:

$$w{=}z*\sqrt[*]{\bar{z}*z}^{\,-1}, \quad \overline{w}{=}\sqrt[*]{\bar{z}*z}^{\,-1}*\bar{z}. \tag{66}$$

It is not hard to obtain

$$[w, \overline{w}]{=}-2\nu*(1{-}w*\overline{w})*(1{-}\overline{w}*w).$$

Hence, we obtain a deformation quantization of the Kähler structure on the Poincaré disk D as a ν-regulated algebra. We denote by $\mathbb{D}_{(\nu)}$ the obtained ν-regulated algebra. The Berezin algebra $\mathbb{D}_{(s)}$ gives an operator representation of $\mathbb{D}_{(\nu)}$, but the positivity of ν is implicitly assumed in advance in the formula of beta functions.

Now consider this in $\mathcal{E}_{2+}(\mathbb{C}^4)$, and let x_0, y_0, x_1, y_1 be a generator system.

$$\mu = e_*^{-2y_0}, \quad \tau = e_*^{y_0}*x_0, \quad \xi = e_*^{y_0}*x_1, \quad \eta = e_*^{y_0}*y_1.$$

Hence, its Heisenberg vacuum representation obtained in § 4 gives an embedding of $\mathcal{H}_2[\mu]$ into $\mathcal{E}_{2+}(\mathbb{C}^4)$ such that $\mu^{-1} \to 1{+}2\hbar s{>}0$ together with other commutation relations.

As above, we set $\zeta{=}\xi{+}i\eta$, $\overline{\zeta}{=}\xi{-}i\eta$. Then, the same calculation as above gives gives exactly the Berezin algebra $\mathbb{D}_{(2s)}$.

References

[1] G. Andrews, R. Askey, R. Roy, SPECIAL FUNCTIONS, Encyclopedia Math, Appl. 71, Cambridge, 2000.

[2] G. S. Agawal, E. Wolf, *Calculus for functions of noncommuting operators and general phase-space method of functions*, Physcal Review D, vol.2, no.10, 1970, 2161-2186.

[3] F.Bayen, M,Flato, C.Fronsdal, A.Lichnerowicz, D.Sternheimer, *Deformation theory and quantization I, II*, Ann. Phys. 111, (1977), 61-151.

[4] F. Berezin, *General concept of quantization,* Comm. Math. Phys. 8 (1975), 153-174.

[5] P. Busch Time-energy uncertainty relation, arXiv:quant-ph/0105049v3, 2007

[6] I.M.Gel'fand, G.E.Shilov, Generalized Functions, 2, Acad. Press, 1968.

[7] L. Hörmander *Fourier Integral operators, I,* Acta Math. 127 (1971), 79-183.

[8] L. Hörmander, *The Weyl calculua of pseudo-differential operators*, Commun. Pure. Appl. Math. 32 (1979) 359-443

[9] N. Hitchin, *Lectures on special Lagrangian submanifolds*, arXiv:math.DG/9907034vl 6Jul, 1999.

[10] N.P. Jewll, *Toeplitz operators on the Bergman space and in several complex variables*, Proc. London Math. Soc. (3) 41,1980 pp193-216.

[11] M. Kontsevitch Deformation quantization of Poisson manifolds, I, qalg/9709040.

[12] H. Omori, INFINITE DIMENSIONAL LIE GROUPS, AMS Translation Monograph 158, 1997.

[13] H. Omori, Y. Maeda, N, Miyazaki, A. Yoshioka, *Deformation quantization of Fréchet-Poisson algebras –Convergence of the Moyal product–*, Math. Phys. Studies. **22** (2000), 233-246.

[14] H. Omori, Y. Maeda, N. Miyazaki, A. Yoshioka, *Deformation quantizations of Fréchet-Poisson algebras of Heisenberg type* , Contemp. Math. **288** (2001), 391-395.

[15] H.Omori, Y.Maeda, N.Miyazaki and A.Yoshioka : *Strange phenomena related to ordering problems in quantizations*, Jour. Lie Theory, Vol 13, No 2, (2003), 481-510.

[16] H.Omori, Y.Maeda, N.Miyazaki and A.Yoshioka : *Deformation of expressions for elements of algebras (I)*, -(Jacobi's theta functions and $*$-exponential functions)- arXiv:1104.2109

[17] H.Omori, Y.Maeda, N.Miyazaki and A.Yoshioka : *Deformation of expressions for elements of algebras (II)*, -(Weyl algebra of $2m$-generators)- arXiv:1105.1218

[18] H.Omori, Y.Maeda, N.Miyazaki and A.Yoshioka : *Deformation of expressions for elements of algebras (III)*, -Generic product formula for $*$-exponentials of quadratic forms- arXiv:1107.2474

[19] H.Omori, Y.Maeda, N.Miyazaki and A.Yoshioka : *Deformation of expressions for elements of algebras (IV)* -Matrix elements and related integrals- arXiv:1109.0082

[20] H.Omori, Y.Maeda, N.Miyazaki and A.Yoshioka : *Deformation of expressions for elements of algebras (V)* -Diagonal matrix calculus and $*$-special functions- arXiv:1111.18062